arXiv:1212.4700v1 [math.PR] 19 Dec 2012

On the convolution powers of complex functions on $\mathbb{Z}$

Evan Randles*[†]
Center for Applied Mathematics
Cornell University

Laurent Saloff-Coste[‡]
Department of Mathematics
Cornell University

Abstract

The local limit theorem describes the behavior of the convolution powers of a probability distribution supported on $\mathbb{Z}$. In this work, we explore the role played by positivity in this classical result and study the convolution powers of the general class of complex valued functions finitely supported on $\mathbb{Z}$. This is discussed as de Forest's problem in the literature and was studied by Schoenberg and Greville. Extending these earlier works and using techniques of Fourier analysis, we establish asymptotic bounds for the sup-norm of the convolution powers and prove extended local limit theorems pertaining to this entire class. As the heat kernel is the attractor of probability distributions on $\mathbb{Z}$, we show that the convolution powers of the general class are similarly attracted to a certain class of analytic functions which includes the Airy function and the heat kernel evaluated at purely imaginary time.

Keywords: Convolution powers, Local limit theorems.

Mathematics Subject Classification: Primary 42A85; Secondary 60F99.

1 Introduction

Let $\phi : \mathbb{Z} \to \mathbb{C}$ be a finitely supported function. We wish to study the convolution powers of ϕ, denoted by $\phi^{(n)}$ and defined iteratively by

$$\phi^{(n)}(x) = \sum_{y\in\mathbb{Z}} \phi^{(n-1)}(x-y)\phi(y)$$

where $\phi^{(1)} = \phi$. This study has been previously motivated by problems in data smoothing and numerical difference schemes for partial differential equa-

---

[*] This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. DGE-0707428

[†] Corresponding author: edr62@cornell.edu

[‡] This material is based upon work supported by the National Science Foundation under Grant No. DMS-1004771

tions [4,7,9,10]. We encourage the reader to see the recent article [2] for background discussion and pointers to the literature.

In the case that the support of ϕ is empty or contains a single point, the convolution powers of ϕ are rather easy to describe. The present article will focus on such functions ϕ with finite support consisting of more than one point; in this case we will say that the support of ϕ is *admissible*. As discussed in [2], when the function ϕ is a probability distribution, i.e., it satisfies $0 \leq \phi(x) \leq 1$ and

$$\sum_{x\in\mathbb{Z}} \phi(x) = 1,$$

the behavior of $\phi^{(n)}$ for large values of n is well-known and is the subject of the local limit theorem (See [6,8]). Our aim is to extend the results of [2] and describe the limiting behavior for the general class of complex valued functions on $\mathbb{Z}$ with admissible support. In particular, we will give bounds on the supremum of $|\phi^{(n)}|$ and prove "generalized" local limit theorems.

As an example, we consider the function $\phi : \mathbb{Z} \to \mathbb{C}$ defined by

$$\phi(0) = \frac{1}{8}(5-2i) \qquad \phi(\pm 1) = \frac{1}{8}(2+i) \qquad \phi(\pm 2) = -\frac{1}{16} \tag{1}$$

and $\phi = 0$ otherwise. The convolution powers $\phi^{(n)}$ for $n = 100, 1000, 10000$ are illustrated in Figures 1 and 2.

We make two crucial observations. First, it appears that the supremum $||\phi^{(n)}||_\infty$ is decaying on the order of $n^{-1/2}$; this is consistent with the classical theory for probability distributions. Second, as n increases, $|\phi^{(n)}(x)|$ appears to be constant for $-100 \leq x \leq 100$. This is in stark contrast to the behavior described by the classical local limit theorem for probability distributions. In the present article, we will prove that there are constants $C, C' > 0$ for which

$$Cn^{-1/2} \leq ||\phi^{(n)}||_\infty \leq C'n^{-1/2}.$$

We will also show that

$$\phi^{(n)}(\lfloor xn^{1/2} \rfloor) = \frac{n^{-1/2}}{\sqrt{4\pi i/8}} e^{-8|x|^2/4i} + o(n^{-1/2})$$

for x in any compact set. Here, $\lfloor \cdot \rfloor$ denotes the greatest integer function. We note that this estimate cannot hold uniformly in x because the modulus of $(4\pi i/8)^{-1/2}\exp(-8|x|^2/4i)$ is a non-zero constant whereas $\phi^{(n)}$ has finite support for each n. For comparison with Figure 2, Figure 3 shows the graph of $\mathrm{Re}((4\pi in/8)^{-1/2}\exp(-8|x|^2/4ni))$ for $n = 100, 1000, 10000$. The above claims will be addressed in Section 8.



Figure 1: $|\phi^{(n)}|$ for $n = 100, 1000, 10000$



Figure 2: $\mathrm{Re}(\phi^{(n)})$ for $n = 100, 1000, 10000$

The Fourier transform is central to the arguments made in this article. We recall its definition: For $\phi : \mathbb{Z} \to \mathbb{C}$, finitely supported, the Fourier transform of



Figure 3: $\mathrm{Re}((4\pi in/8)^{-1/2}\exp(-8|x|^2/4ni))$ for $n = 100, 1000, 10000$

ϕ is

$$\hat{\phi}(\xi) = \sum_{x\in\mathbb{Z}} \phi(x)e^{ix\xi}. \tag{2}$$

Our first main result is illustrated in the following theorem.

Theorem 1. *Let $\phi : \mathbb{Z} \to \mathbb{C}$ have admissible support and let $A = \sup_\xi |\hat{\phi}(\xi)|$. Then there is a natural number $m \geq 2$, and positive constants C and C' such that*

$$Cn^{-1/m} \leq A^{-n}||\phi^{(n)}||_\infty \leq C'n^{-1/m} \tag{3}$$

for all natural numbers n.

In the classical local limit theorem, the convolution powers of a probability distribution are approximated by the heat kernel, an analytic function. In the present more general setting, the convolution powers, $\phi^{(n)}$, are analogously approximated by certain analytic functions. We define these now. Let $m \geq 2$ be a natural number and β be a non-zero complex number such that $\mathrm{Re}(\beta) \geq 0$. We define

$$H_m^\beta(x) = \frac{1}{2\pi}\int_{\mathbb{R}} e^{-ixu}e^{-\beta u^m}du \tag{4}$$

when the above integral converges. When $\mathrm{Re}(\beta) > 0$ and m is an even natural number it is easy to see that

$$|e^{-ixu}e^{-\beta u^m}| = e^{-\mathrm{Re}(\beta)u^m} \in L^1(\mathbb{R})$$

whence the convergence is uniform in x. In this case, H_m^β is equivalently defined by its Fourier transform, $e^{-\beta u^m}$, which exists in the sense of Schwartz functions. In the case that $\mathrm{Re}(\beta) = 0$, it isn't immediately clear for which values of m or in what sense the integral in (4) will converge. It will be shown that for $m > 2$ the convergence is uniform in x and when $m = 2$ the convergence is uniform for x in any compact set. This is the subject of Proposition 2.

Remark 1. *In the case that* $\mathrm{Re}(\beta) > 0$ *the function* H_m^β *and the function* $H_{m,b}$ *used in Theorem 2.3 of [2] and defined by its Fourier transform,* $\hat{H}_{m,b}(\xi) = e^{-(1+ib)\xi^m}$, *are connected via the relation*

$$H_{m,\frac{\mathrm{Im}(\beta)}{\mathrm{Re}(\beta)}}\left(\frac{x}{(\mathrm{Re}(\beta))^{1/m}}\right) = (\mathrm{Re}(\beta))^{1/m} H_m^\beta(x).$$

Let us look at some specific instances of H_m^β. When $m = 2$ the function

$$H_m^\beta(x) = \frac{1}{\sqrt{4\pi\beta}} e^{-\frac{|x|^2}{4\beta}} \tag{5}$$

is the heat kernel evaluated at complex time β. That is, $H_m^\beta(x)$ is the kernel of the operator $e^{\beta\Delta}$ where Δ is the unique self-adjoint extension of $(d/dx)^2$ originally defined on smooth compactly supported functions on $\mathbb{R}$. In the case that $m \geq 2$ and $\beta = i/m$, $H_m^\beta = H_m^{i/m}$ satisfies the ordinary differential equation

$$\frac{d^{m-1}y}{dx^{m-1}} + i^{m-1}xy = 0.$$

When $m = 3$ this is Airy's equation and $H_3^{i/3}(x)$ is the famous Airy function, $\mathrm{Ai}(x)$. The graphs of $\mathrm{Re}(H_m^i(x))$ for $m = 2, 3, 4, 5$ are illustrated in Figure 4.

In addition to the convergence of the defining integrals, we are interested in studying other properties of the function $H_m^\beta(x)$. Such properties will allow us to easily deduce information about the convolution powers of ϕ. In particular, we will study the asymptotic behavior of $H_m^\beta(x)$. When $\mathrm{Re}(\beta) > 0$ and m is an even natural number it is easily shown that

$$|H_m^\beta(x)| \leq C \exp(-B|x|^{\frac{m}{m-1}}) \tag{6}$$

for all real numbers x where $C, B > 0$ (See [3]). In the case where $\mathrm{Re}(\beta) = 0$ the situation is quite different. When $m = 2$, (5) implies that $|H_m^\beta(x)|$ is constant for all $x \in \mathbb{R}$. When $m > 2$ we will show that

$$|H_m^\beta(x)| \leq \frac{A}{|x|^{\frac{(m-2)}{2(m-1)}}} + \frac{B}{|x|}$$

for all real numbers x where $A, B > 0$. It is interesting to note that when $m = 3$ the above estimate is a statement about the asymptotic behavior of the Airy function and is usually found using the method of steepest descent. In this case



Figure 4: $\mathrm{Re}(H_m^i(x))$ for $m = 2, 3, 4, 5$

our estimate is sharp.

The latter two main results of this article, Theorems 2 and 3, are both stated under the assumption that $\sup_\xi |\hat{\phi}(\xi)| = 1$; this can always be arranged by replacing ϕ by $A\phi$ for an appropriate constant $A > 0$. Without further ado, our main local limit theorem is as follows:

Theorem 2. *Let $\phi : \mathbb{Z} \to \mathbb{C}$ have admissible support and be such that $\sup_\xi |\hat{\phi}(\xi)| = 1$. Then there is a natural number $m \geq 2$ and a finite collection of real numbers $\alpha_1, \alpha_2, \ldots, \alpha_Q$ such that for each α_q in the above collection, there are complex numbers $\beta_1, \beta_2, \ldots, \beta_{R_q}$ and points $\xi_1, \xi_2, \ldots, \xi_{R_q} \in (-\pi, \pi]$ such that*

$$\phi^{(n)}(\lfloor \alpha_q n + x n^{1/m} \rfloor) = \sum_{l=1}^{R_q} n^{-1/m} e^{-i\lfloor x n^{1/m} + \alpha_q n \rfloor \xi_l} \hat{\phi}(\xi_l)^m H_m^{\beta_l}(x) + o(n^{-1/m}) \tag{7}$$

uniformly for x in any compact set.

As we will show, when ϕ satisfies one mild additional hypothesis we can say a great deal more about the convolution powers of ϕ. This is seen in the following theorem. In specific cases the theorem recaptures results originally proved by Schoenberg in [7], Greville in [4] and Thomée in [10] (See also [2]).

Theorem 3. *Let $\phi : \mathbb{Z} \to \mathbb{C}$ have admissible support and be such that* $\sup_\xi |\hat{\phi}(\xi)| = 1$. *Additionally assume that for every $\rho \in (-\pi, \pi]$ for which $|\hat{\phi}(\rho)| = 1$ we have*

$$\frac{d^2}{d\xi^2} \log(\hat{\phi}(\xi))\Big|_\rho \neq i\tau \tag{8}$$

for any non-zero real number τ. Then there exist a natural number $m \geq 2$, a compact set K, a finite number of points $\xi_1, \xi_2, \ldots, \xi_Q$, real numbers $\alpha_1, \alpha_2, \ldots, \alpha_Q$ and complex numbers $\beta_1, \beta_2, \ldots, \beta_Q$ such that $||\phi^{(n)}||_\infty$ is attained on

$$\bigcup_{q=1}^{Q} (\alpha_q n + K n^{1/m}) \tag{9}$$

and

$$\phi^{(n)}(x) = \sum_{q=1}^{Q} n^{-1/m} e^{-ix\xi_q} \hat{\phi}(\xi_q)^n H_m^{\beta_q}\left(\frac{x - \alpha_q n}{n^{1/m}}\right) + o(n^{-1/m}) \tag{10}$$

uniformly in $\mathbb{Z}$.

Remark 2. *The α's and β's in the statements of Theorems 2 and 3 are computed via the Taylor expansion of* $\log(\hat{\phi}(\xi))$ *about points ξ_q for which $|\hat{\phi}(\xi_q)| = 1$. This is the focus of Section 2. These constants play the analogous roles of the mean and first non-vanishing moment of order $m > 1$ for probability distributions. Our more general setting allows for the existence of multiple such constants (See Theorem 1 in Section 2.7 of [8]).*

The paper is organized as follows: In Section 2 we study the local behavior of the Fourier transform of ϕ. Section 3 develops the machinery involving oscillatory integrals that will be used in proving the local limit theorems. As an easy consequence of this we finish the section by proving the upper estimate of Theorem 1; this is Theorem 4. In section 4, we focus on the functions H_m^β where we address convergence, analyticity and asymptotic bounds. The local limit theorems of Theorems 2 and 3 are proven in Section 5. In Section 6, we complete the proof of Theorem 1 and in Section 7 the first conclusion of Theorem 2 is proven. Section 8 gives examples and addresses a general situation previously treated in [2].

2 Local behavior of $\hat{\phi}$

In this section we study the local behavior of $\hat{\phi}$ at points for which the supremum of $|\hat{\phi}|$ is attained. This will be seen to completely determine the limiting behavior of the convolution powers of ϕ. In particular, the Taylor series of the logarithm of $\hat{\phi}$ will describe the local behavior of $\hat{\phi}$, as well as give the constants m, $\alpha_1, \alpha_2, \ldots, \alpha_Q$ and $\beta_1, \beta_2, \ldots, \beta_R$ used in the statements of Theorems 2 and 3. We proceed by making some simple observations about (2) under the assumptions that ϕ has admissible support and $\sup_\xi |\hat{\phi}(\xi)| = 1$.

Our first observation concerns the number of points at which $|\hat{\phi}(\xi)| = 1$. It follows from the assumptions above that $|\hat{\phi}|^2$ is a non-constant trigonometric polynomial and so $|\hat{\phi}|$ is not constant. From here we observe that $\hat{\phi}$ can only satisfy $|\hat{\phi}(\xi)| = 1$ at a finite number of points in $(-\pi, \pi]$; a simple accumulation-point argument shows the necessity of this fact. We now observe that $\hat{\phi}$ is a finite linear combination of exponentials and is therefore analytic. We use this observation to study the local behavior of $\hat{\phi}(\xi)$ about any point ξ_0 for which $|\hat{\phi}(\xi_0)| = 1$. To this end we consider

$$\eta(\xi) = \log\left(\frac{\hat{\phi}(\xi + \xi_0)}{\hat{\phi}(\xi_0)}\right) \tag{11}$$

where log is taken to be the principle branch of logarithm and we allow the variable ξ to be complex, for the time being. It follows from our remarks above that η is analytic on an open neighborhood of 0 and we can therefore consider its convergent Taylor series

$$\eta(\xi) = \sum_{l=0}^{\infty} a_l \xi^l$$

on this neighborhood. The limiting behavior of the convolution powers of ϕ will be in some sense characterized by the first few non-zero terms of this series.

The requirement that $|\hat{\phi}(\xi)| \leq 1$ imposes conditions on the Taylor expansion for η as follows: We consider the collection $\{a_l\}_{l=1}^{\infty}$ of coefficients of the series and let $k = \min\{l : \mathrm{Re}(a_l) \neq 0\}$, which exists, for otherwise $|\hat{\phi}|$ would be constant. We claim that k is even and $\mathrm{Re}(a_k) < 0$. To see this we observe that by only considering real values of ξ we can find a neighborhood of 0 on which

$$e^{C\xi^k} \leq |\hat{\phi}(\xi + \xi_0)| = |\hat{\phi}(\xi_0) e^{\eta(\xi)}|$$

and where C is a real constant sharing the same sign as $\mathrm{Re}(a_k)$. If it is the case that $\mathrm{Re}(a_k) > 0$ or k is odd we have that $|\hat{\phi}(\xi + \xi_0)| > 1$ for some ξ which cannot be true. We will summarize the above arguments shortly. First we give the following convenient definition, originally motivated by Thomée [10].

Definition 1. *Let $\nu : \mathbb{R} \to \mathbb{C}$ be analytic on a neighborhood of a point ξ_0 such that $|\nu(\xi_0)| = 1$. Let $\eta : \mathbb{R} \to \mathbb{C}$ be defined by*

$$\eta(\xi) = \log\left(\frac{\nu(\xi + \xi_0)}{\nu(\xi_0)}\right). \tag{12}$$

1. *We say that ξ_0 is a point of type 1 and of order m for ν if the Taylor expansion for* (12) *yields an even integer m, a real number α and a complex number β with* $\mathrm{Re}(\beta) > 0$ *such that*

$$\eta(\xi) = i\alpha\xi - \beta\xi^m + \sum_{l=m+1}^{\infty} a_l \xi^l \tag{13}$$

on a neighborhood of 0*. We will write* $\xi_0 \in (1; m)$*.*

2. *We say that* ξ_0 *is a point of type 2 and of order* m *for* ν *if the Taylor expansion for* (12) *yields* $m, k, \alpha, \gamma, p(\xi)$ *where* m *and* k *are natural numbers with* k *even and* $1 < m < k$*,* α *and* γ *are real numbers with* $\gamma > 0$ *and,* $p(\xi)$ *is a real polynomial with* $p(0) \neq 0$ *such that*

$$\eta(\xi) = i\alpha\xi - i\xi^m p(\xi) - \gamma\xi^k + \sum_{l=k+1}^{\infty} a_l \xi^l. \tag{14}$$

on a neighborhood of 0*. We will set* $\beta = ip(0)$ *and write* $\xi_0 \in (2; m)$*.*

In either case we will refer to the constant α *as the drift constant for* ξ_0*.*

Using the above definition, the arguments which preceded it are summarized in the following proposition.

Proposition 1. *Let* $\phi : \mathbb{Z} \to \mathbb{C}$ *have admissible support. Suppose that the Fourier transform of* ϕ *satisfies* $\sup|\hat{\phi}(\xi)| = 1$*. Then there are only a finite number of points,* $\xi_1, \xi_2, \ldots, \xi_R \in (-\pi, \pi]$ *such that* $|\hat{\phi}(\xi_q)| = 1$*. For each of these points,* ξ_q*, we have either* $\xi_q \in (1; m)$ *or* $\xi_q \in (2; m)$*.*

The following Lemma will be seen to govern the limiting behavior of the convolution powers of ϕ.

Lemma 1. *Let* $\nu : \mathbb{R} \to \mathbb{C}$ *be analytic on a neighborhood of a point* ξ_0 *such that* $|\nu(\xi_0)| = 1$*.*
1. *If* $\xi_0 \in (1; m)$*, then*

$$\nu(\xi_0 + \xi) = \nu(\xi_0) \exp(i\alpha\xi - \beta\xi^m(1 + o(1)) \text{ as } \xi \to 0 \tag{15}$$

and

$$|\nu(\xi + \xi_0)| \leq e^{-C\xi^m} \tag{16}$$

for some constant $C > 0$ *and for* ξ *in a neighborhood of* 0*. Here* α *and* β *are given by Definition 1.*

2. *If* $\xi_0 \in (2; m)$*, then*

$$\nu(\xi_0 + \xi) = \nu(\xi_0) \exp(i\alpha\xi - i\xi^m p(\xi) - \gamma\xi^k(1 + o(1)) \text{ as } \xi \to 0 \tag{17}$$

and the function

$$g(u) = [\nu(\xi_0)^{-1}\nu(\xi + \xi_0) \exp(-i\alpha\xi + i\xi^m p(\xi))]$$

satisfies

$$|g(\xi)| \leq e^{-C\xi^k} \tag{18}$$

and

$$|g'(\xi)| \leq D|\xi|^{k-1} e^{-C\xi^k} \tag{19}$$

for some constants $C, D > 0$ *and for* ξ_0 *in a neighborhood of* 0*. Here* $k, \alpha, p(\xi)$ *and* γ *are given by Definition 1.*

Proof. As an immediate consequence of our definitions we have

$$\nu(\xi + \xi_0) = \nu(\xi_0)e^{\eta(\xi)}$$

where η is defined by (12). In the case that $\xi_0 \in (1; m)$, (15) and (16) are immediate from (13) and the fact that the series $\sum_{l=m+1} a_l \xi^l$ converges uniformly on a neighborhood of 0.

In the case that $\xi_0 \in (2; m)$, the justification of the estimates (17) and (18) follows similarly. For the last conclusion we observe that

$$\begin{aligned} g'(\xi) &= \frac{d}{d\xi}\exp(-i\alpha\xi + i\xi^m p(\xi))e^{\eta(\xi)} \\ &= \frac{d}{d\xi}\exp\left(-\gamma\xi^k + \sum_{l=k+1}^{\infty} a_l\xi^l\right) \\ &= \left(-\gamma k\xi^{k-1} + \sum_{l=k+1}^{\infty} a_l l\xi^{l-1}\right) g(\xi) \end{aligned}$$

on a neighborhood of 0. The inequality (19) now follows without trouble. □

3 The upper bound

The goal of this section is to establish the upper bound of Theorem 1. In the process we will prove a series of technical lemmas involving oscillatory integrals of the form

$$\int_a^b g(\xi)e^{if(\xi)}d\xi$$

which will be used often throughout the remainder of the paper. Many of the arguments within are based on the same or slightly less general arguments made by Greville [4], Tomée [10] and, not surprisingly, van der Corput.

Lemma 2. *Let $h \in L^1([a,b])$ and $g \in \mathcal{C}^1([a,b])$ be complex valued. Then for any M such that*

$$\left|\int_a^x h(u)du\right| \leq M$$

for all $x \in [a,b]$ we have

$$\left|\int_a^b g(u)h(u)du\right| \leq M\left(||g||_\infty + ||g'||_1\right).$$

Proof. For $h \in L^1([a,b])$, the function

$$f(x) = \int_b^x h(u)du$$

is absolutely continuous and $f'(x) = h(x)$ almost everywhere. Moreover our hypothesis guarantees that $|f(x)| \leq M$ for all $x \in [a,b]$. Integrating by parts yields

$$\int_a^b g(u)h(u)du = [g(u)f(u)]_a^b - \int_a^b g'(u)f(u)du$$

and therefore

$$\begin{aligned} \left| \int_a^b g(u)h(u)du \right| &\leq |f(b)g(b)| + 0 + \int_a^b |f(u)||g'(u)|du \\ &\leq M||g||_\infty + M||g'||_1. \end{aligned}$$

□

The following two lemmas are due to van der Corput, their proofs can be found in [10] and [1]. The proof of the former is a nice application of the second mean value theorem for integrals. Its validity can also be seen using alternating series, See Lemma 3 of [4].The proof of the latter is relatively simple but involves checking several cases.

Lemma 3. *Let $f \in \mathcal{C}^1([a,b])$ be real valued and suppose that f' is a monotonic function such that $f'(x) \neq 0$ for all $x \in [a,b]$. Then,*

$$\left| \int_a^b e^{if(u)}du \right| \leq \frac{4}{\lambda} \tag{20}$$

where

$$\lambda = \inf_{x \in [a,b]} |f'(x)|. \tag{21}$$

Lemma 4. *Let $f \in \mathcal{C}^2([a,b])$ be real valued and suppose that $f''(x) \neq 0$ for all $x \in [a,b]$. Then*

$$\left| \int_a^b e^{if(u)}du \right| \leq \frac{8}{\sqrt{\rho}}$$

where

$$\rho = \inf_{x \in [a,b]} |f''(x)|.$$

Lemma 5. *Let $g \in \mathcal{C}^1([a,b])$ be complex valued and let $f \in C^2([a,b])$ be real valued and such that $f''(x) \neq 0$ for all $x \in [a,b]$. Then*

$$\left| \int_a^b g(u)e^{if(u)}du \right| \leq \min\left\{ \frac{4}{\lambda}, \frac{8}{\sqrt{\rho}} \right\} (||g||_\infty + ||g'||_1)$$

where $\lambda = \inf_{x \in [a,b]} |f'(x)|$ and $\rho = \inf_{x \in [a,b]} |f''(x)|$.

Proof. Combining the results of Lemmas 3 and 4 show

$$\left|\int_a^x e^{if(u)}du\right| \leq \min\left\{\frac{4}{\lambda}, \frac{8}{\sqrt{\rho}}\right\}$$

for any $x \in [a,b]$. We remark that $4/\lambda$ will only contribute to the upper bound provided f' is never zero and in which case the application of Lemma 3 is justified. Setting $h(u) = e^{if(u)}$ we note that the functions g and h are the subject of Lemma 2. The result now follows immediately from Lemma 2. ☐

Lemma 6. *Let $\nu : \mathbb{R} \to \mathbb{C}$ be analytic on a neighborhood of ξ_0 where $|\nu(\xi_0)| = 1$. If ξ_0 is a point of order $m \geq 2$ for ν, then there is $\delta > 0$ such that*

$$\frac{1}{2\pi}\int_{|\xi-\xi_0|\leq\delta} \nu(\xi)^n e^{-ix\xi}d\xi = O(n^{-1/m})$$

where the limit is uniform in $x \in \mathbb{R}$.

Proof. Let us first assume that $\xi_0 \in (1;m)$. Our hypothesis guarantees that m is even and by Lemma 1 there are $C > 0$ and $\delta > 0$ such that

$$|\nu(\xi+\xi_0)| \leq e^{-C\xi^m}$$

for all $-\delta \leq \xi \leq \delta$. Therefore

$$\begin{aligned}\left|\frac{1}{2\pi}\int_{|\xi-\xi_0|\leq\delta} \nu(\xi)^n e^{-ix\xi}d\xi\right| &\leq \int_{-\delta}^{\delta} |\nu(\xi+\xi_0)|^n d\xi \\ &\leq \int_{\mathbb{R}} e^{-nC\xi^m}d\xi \\ &\leq \frac{M}{n^{1/m}}.\end{aligned}$$

In the second case we assume that $\xi_0 \in (2;m)$. We set

$$g(\xi) = \left[\nu(\xi_0)^{-1}\nu(\xi+\xi_0)\exp(-i\alpha\xi + i\xi^m p(\xi))\right]$$

and

$$f_n(\xi, x) = (n\alpha - x)\xi - n\xi^m p(\xi).$$

We note that f_n is real valued. Appealing to Lemma 1, let $\delta > 0$ be chosen so that the estimates (18) and (19) hold for all $\xi \in [-\delta, \delta]$. Upon changing variables of integration and using the fact that $|\nu(\xi_0)| = 1$ we may write

$$\left|\frac{1}{2\pi}\int_{|\xi-\xi_0|\leq\delta} \nu(\xi)^n e^{-ix\xi}d\xi\right| \leq \sum_{j=1}^{3}\left|\int_{I_j} g(\xi)^n e^{if_n(\xi,x)}d\xi\right|$$

where $I_1 = [-\delta, -n^{1/m}]$, $I_2 = (-n^{1/m}, n^{1/m})$ and $I_3 = [n^{1/m}, \delta]$. On the interval I_2, $|g(\xi)| \leq 1$ by (18) and therefore

$$\left|\int_{I_2} g(\xi)^n e^{if_n(\xi,x)}d\xi\right| \leq \frac{2}{n^{1/m}}.$$

We now consider the integral over I_1 to which we will apply Lemma 5. First observe that the regularity requirements of Lemma 5 for f_n and g^n are met. Differentiating f_n twice with respect to ξ gives

$$\partial_\xi^2 f_n(\xi,x) = -n\frac{d^2}{d\xi^2}\xi^m p(\xi),$$

which is independent of x. Using the fact that $\xi^m p(\xi)$ is a polynomial with m being the smallest power of its terms, we may further restrict $\delta > 0$ so that

$$C^2|\xi|^{m-2} \leq \left|\frac{d^2}{d\xi^2}\xi^m p(\xi)\right| \tag{22}$$

for some $C > 0$ and for all $\xi \in [-\delta, \delta]$. Consequently $|\partial_\xi^2 f_n(\xi,x)| > 0$ for all $\xi \in I_1$ and $x \in \mathbb{R}$. Appealing to Lemma 4 we have

$$\left|\int_{I_1} g(\xi)^n e^{if_n(\xi,x)} d\xi\right| \leq \frac{8}{\sqrt{\lambda}}(||g^n||_\infty + ||ng' g^{n-1}||_1) \tag{23}$$

where $\lambda = \inf_{\xi\in I_1} |\partial_\xi^2 f_n(\xi,x)|$ and we have used the power rule for the derivative of g^n. Using (22) and recalling that $m \geq 2$ we observe that

$$Cn^{1/m} = \sqrt{C^2 n| - n^{-1/m}|^{m-2}} \leq \sqrt{\inf_{I_1} C^2 n|\xi|^{m-2}} \leq \sqrt{\lambda}.$$

Now by (18) and (19) of Lemma 1 we have $||g^n|| \leq 1$ and

$$\begin{aligned} ||ng' g^{n-1}||_1 &= n\int_{I_1} |g'(\xi) g(\xi)^n| d\xi \\ &\leq n\int_{I_1} D|\xi|^{k-1} e^{-nC\xi^k} d\xi \\ &\leq \int_{\mathbb{R}} D|u|^{k-1} e^{-Cu^k} du = M < \infty. \end{aligned}$$

Inserting the above estimates into (23) gives

$$\left|\int_{I_1} g(\xi)^n e^{if_n(\xi,x)} d\xi\right| \leq \frac{8(1+M)}{Cn^{1/m}} = \frac{K_1}{n^{1/m}}.$$

A similar calculation shows that

$$\left|\int_{I_3} g_n(\xi) e^{if_n(\xi,x)} d\xi\right| \leq \frac{K_2}{n^{1/m}}$$

for some $K_2 > 0$. Putting these estimates together gives

$$\left|\frac{1}{2\pi}\int_{|\xi-\xi_0|\leq\delta} \nu(\xi)^n e^{-ix\xi} d\xi\right| \leq \frac{K_1}{n^{1/m}} + \frac{2}{n^{1/m}} + \frac{K_2}{n^{1/m}},$$

our desired inequality. $\square$

Theorem 4. *Let* $\phi : \mathbb{Z} \to \mathbb{C}$ *have admissible support and let* $A = \sup_\xi |\hat{\phi}(\xi)|$. *Then there is a natural number* $m \geq 2$ *and a real number* $C' > 0$ *such that*

$$A^{-n}||\phi^{(n)}||_\infty \leq C' n^{-1/m} \tag{24}$$

for all natural numbers n.

Proof. For any natural number n we have the identity

$$\phi^{(n)}(x) = \frac{1}{2\pi}\int_{-\pi}^{\pi} \hat{\phi}(\xi)^n e^{-ix\xi} d\xi. \tag{25}$$

By inspecting (25) we remark that it suffices to prove the theorem in the case that $A = \sup_\xi |\hat{\phi}(\xi)| = 1$, for otherwise one simply multiplies (25) by A^{-n}. Using Proposition 1 let $\xi_1, \xi_2, \ldots, \xi_Q$ be the set of points at which $|\hat{\phi}(\xi)| = 1$. By the proposition we know that for each $q = 1, 2, \ldots, Q$ there is a natural number $m_q \geq 2$ such that ξ_q is a point of order m_q for $\hat{\phi}$. Appealing to Lemma 6 for each ξ_q we choose $\delta_q > 0$ for which the conclusion of the lemma holds. Set $m = \max m_q$, $I_q = [\xi_q - \delta_q, \xi_q + \delta_q]$, $J = [-\pi, \pi) \setminus \cup_q I_q$ and $s = \sup_\xi |\hat{\phi}(\xi)| < 1$. We write

$$\begin{aligned}
|\phi^{(n)}(x)| &= \left| \sum_{q=1}^{Q} \frac{1}{2\pi}\int_{I_q} \hat{\phi}(\xi)^n e^{-ix\xi} d\xi + \frac{1}{2\pi}\int_J \hat{\phi}(\xi)^n e^{-ix\xi} d\xi \right| \\
&\leq \sum_{q=1}^{Q} \left| \frac{1}{2\pi}\int_{I_q} \hat{\phi}(\xi)^n e^{-ix\xi} d\xi \right| + \frac{1}{2\pi}\int_J |\hat{\phi}(\xi)|^n d\xi \\
&\leq \sum_{q=1}^{Q} \left| \frac{1}{2\pi}\int_{I_q} \hat{\phi}(\xi)^n e^{-ix\xi} d\xi \right| + s^n.
\end{aligned}$$

Using Lemma 6 we conclude that for every $x \in \mathbb{R}$

$$\begin{aligned}
|\phi^{(n)}(x)| &\leq \sum_{q=1}^{Q} \frac{K_q}{n^{1/m_q}} + s^n \\
&\leq \frac{K}{n^{1/m}} + s^n
\end{aligned}$$

from which the result follows. □

4 The attractors H_m^β

In this section we study the functions H_m^β defined by (4). We will first show that the integral defining H_m^β converges in the senses indicated in the introduction.

Proposition 2. *Let* $m \geq 2$ *be a natural number and let* β *be a non-zero complex number such that* $\mathrm{Re}(\beta) \geq 0$.

1. *If m is even and $\operatorname{Re}\beta > 0$ then the integral defining $H_m^\beta(x)$ in* (4) *converges uniformly for all $x \in \mathbb{R}$.*
2. *If $m > 0$ and $\operatorname{Re}(\beta) = 0$ then the integral defining $H_m^\beta(x)$ converges uniformly for all $x \in \mathbb{R}$.*
3. *If $m = 2$ and $\operatorname{Re}(\beta) = 0$ then for any compact set $K \subseteq \mathbb{R}$ the integral defining $H_m^\beta(x)$ converges uniformly for all $x \in K$.*

Proof. For item 1 there is nothing to prove, the result follows from the classical theory of Fourier transforms. For items 2 and 3 we may let τ be the non-zero real number such that $\beta = i\tau$ and set $f(u,x) = -xu - \tau u^m$. Our job is to show that the integral

$$\int_{\mathbb{R}} e^{if(x,u)} du$$

converges in the senses indicated for $m > 2$ and $m = 2$ respectively.

We first consider the case where $m > 2$. Let $\epsilon > 0$ and choose M sufficiently large so that

$$\frac{8}{\sqrt{|\tau| m(m-1)M^{m-2}}} \leq \epsilon. \tag{26}$$

Observe that for any real numbers a and b such that $M < a \leq b$ or $a \leq b < -M$,

$$|\tau| m(m-1)M^{m-2} < \inf_{u \in [a,b]} |\partial_u^2 f(x,u)| = \lambda.$$

We now apply Lemma 4 and conclude that

$$\left| \int_a^b e^{-ixu - \beta u^m} du \right| = \left| \int_a^b e^{if(x,u)} du \right| \leq \frac{8}{\sqrt{\lambda}} < \epsilon$$

for all $x \in \mathbb{R}$ and for all $a \leq b$ such that the distance from the interval $[a,b]$ to 0 is more than M. The Cauchy criterion for indefinite integrals guarantees that the integrals

$$\int_0^\infty e^{-ixu - \beta u^m} du \text{ and } \int_{-\infty}^0 e^{-ixu - \beta u^m} du$$

converge uniformly for all $x \in \mathbb{R}$. This proves item 2.

Let us now assume that $m = 2$. We remark that the above argument fails in this case because the $\partial_u^2 f$ is a non-zero constant. We will consequently need to use $\partial_u f$, which depends on both u and x, to bound our integrals. Let $\epsilon > 0$ and let $K \subseteq \mathbb{R}$ be a compact set. We choose $M > 0$ so that

$$\frac{4}{|2\tau M + x|} < \epsilon$$

for all $x \in K$. By applying Lemma 3 and making an argument analogous to that given in the previous case we conclude that

$$\left| \int_a^b e^{-ixu - \beta u^2} du \right| = \left| \int_a^b e^{if(x,u)} du \right| < \epsilon$$

for all $x \in K$ and for all $a \leq b$ such that the distance from the interval $[a, b]$ to 0 is more than M. Again an application of the Cauchy criterion gives the desired result. ☐

Proposition 3. *Let β be non-zero and purely imaginary. Then for any natural number $m > 2$ there exists positive constants A, B such that*

$$|H_m^\beta(x)| \leq \frac{A}{|x|^{\frac{m-2}{2(m-1)}}} + \frac{B}{|x|} \tag{27}$$

for all $x \in \mathbb{R}$.

Proof. Let $\beta = i\tau$ where τ is a non-zero real number. By Proposition 2 we know that for any $\epsilon > 0$ there is $M > 0$ such that

$$|H_m^\beta(x)| \leq \frac{1}{2\pi} 2\epsilon + \frac{1}{2\pi} \left| \int_{-M}^{M} e^{-i(xu+\tau u^m)} du \right| \tag{28}$$

for all $x \in \mathbb{R}$. We have actually shown more. The proof of Proposition 2, in particular (28), shows that when M satisfies $(8/\epsilon)^2 = |\tau| m(m-1)M^{m-2}$ the estimate above holds for all $x \in \mathbb{R}$. We apply Lemma 3 to the above integral and conclude that for any $x \in \mathbb{R}$ and $\epsilon > 0$

$$|H_m^\beta(x)| \leq \epsilon + \frac{1}{\lambda} \tag{29}$$

where $\lambda = \inf_{u \in [-M, M]} |x + \tau m u^{m-1}|$. A careful inspection, considering all possible values of m, τ and M, shows that

$$\big| |x| - |\tau| m M^{m-1} \big| \leq \lambda$$

provided $|x| \geq |\tau| m M^{m-1}$, a condition that is trivially satisfied when $|x| = 2|\tau| m M^{m-1}$. Consequently for any $x \in \mathbb{R}$

$$|H_m^\beta(x)| \leq \epsilon + \frac{1}{|\tau| m M^{m-1}}$$

subject to the constraints $|x| = 2|\tau| m M^{m-1}$ and $(8/\epsilon)^2 = |\tau| m(m-1)M^{m-2}$. It now follows that for every $x \in \mathbb{R}$

$$|H_m^\beta(x)| \leq \left(\frac{2^{\frac{m-2}{m-1}}}{(m-1)(|\tau| m)^{1/(m-1)}} \right)^{1/2} \frac{8}{|x|^{\frac{m-2}{2(m-1)}}} + \frac{2}{|x|}, \tag{30}$$

our desired result. ☐

It is not surprising that the functions H_m^β are analytic (and not identically zero) for all values of m and β considered above. When m is even and $\mathrm{Re}(\beta) > 0$, analyticity follows from the classical theory of Fourier transforms. In the case that m is odd and β is purely imaginary the result was proved by R. Hersch in [5]

(See Theorem 4 therein). The proof involves changing the contour of integration from $\mathbb{R}$ to a pair of rays on which the integrand is absolutely integrable. In the case that m is even and β is purely imaginary, Hersh's argument pushes through with very little modification. We will therefore not provide a proof for this final case and encourage the reader to see [5] for the essential details. The result is as follows:

Proposition 4. *Let $m \geq 2$ be a natural number and let β be a non-zero complex number with $\operatorname{Re}(\beta) \geq 0$. If $\operatorname{Re}(\beta) > 0$ additionally assume that m is even. Then H_m^β is analytic and not identically zero.*

5 Local limits

Lemma 7. *Let $\nu : \mathbb{R} \to \mathbb{C}$ be analytic on a neighborhood of a point ξ_0 such that $|\nu(\xi_0)| = 1$. Assume that $\xi_0 \in (1; m)$ with associated constants α and β. Then for all $\epsilon > 0$ there is a $\delta > 0$ and a natural number N such that*

$$\left| \frac{n^{1/m}}{2\pi} \int_{|\xi-\xi_0|\leq\delta} \nu(\xi)^n e^{-ix\xi} d\xi - e^{-ix\xi_0} \nu(\xi_0)^n H_m^\beta \left(\frac{x-\alpha n}{n^{1/m}} \right) \right| < \epsilon \tag{31}$$

for all $n > N$ and for all real numbers x.

Proof. Let $\epsilon > 0$ and set,

$$y_n = (x - \alpha n) n^{-1/m}$$

and

$$g(u) = \left[\nu(\xi_0)^{-1} e^{-i\alpha u n^{-1/m}} \nu(\xi_0 + u n^{-1/m}) \right].$$

Upon changing variables of integration we have

$$\begin{aligned} &\frac{n^{1/m}}{2\pi} \int_{|\xi-\xi_0|\leq\delta} \nu(\xi)^n e^{-ix\xi} d\xi \\ &= \frac{e^{-ix\xi_0}\nu(\xi_0)^n}{2\pi} \int_{|u|\leq\delta n^{1/m}} \left[\nu(\xi_0)^{-1} e^{-i\alpha u n^{-1/m}} \nu(\xi_0 + u n^{-1/m}) \right]^n e^{-iu\frac{x-\alpha n}{n^{1/m}}} du \\ &\qquad = \frac{e^{-ix\xi_0}\nu(\xi_0)^n}{2\pi} \int_{|u|\leq\delta n^{1/m}} g(u)^n e^{-iuy_n} du. \end{aligned}$$

Comparing the above integral with $e^{-ix\xi_0}\nu(\xi_0)^n H_m^\beta(y_n)$ gives

$$\begin{aligned}
&\left|\frac{n^{1/m}}{2\pi}\int_{|\xi-\xi_0|\le\delta}\nu(\xi)^n e^{-ix\xi}d\xi - e^{-ix\xi_0}\nu(\xi_0)^n H_m^\beta(y_n)\right| \\
&\qquad\qquad = \left|\frac{e^{-ix\xi_0}\nu(\xi_0)^n}{2\pi}\int_{|u|\le\delta n^{1/m}}[g(u)^n - e^{-\beta u^m}]e^{-iuy_n}du\right| \\
&\qquad\qquad\qquad + \left|\frac{e^{-ix\xi_0}\nu(\xi_0)^n}{2\pi}\int_{|u|>\delta n^{1/m}} e^{-\beta u^m}e^{-iuy_n}du\right| \\
&\qquad\qquad \le \int_{|u|\le\delta n^{1/m}}|g(u)^n - e^{-\beta u^m}|du + \int_{|u|>\delta n^{1/m}} e^{-\operatorname{Re}(\beta)u^m}du \\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad = \mathcal{I}_1 + \mathcal{I}_2.
\end{aligned}$$

Now we estimate the integrals $\mathcal{I}_1$ and $\mathcal{I}_2$. For $\mathcal{I}_1$ It follows from (15) and our definition of g that for any $\eta > 0$ we may choose $\delta > 0$ for which

$$|g(u)^n - e^{-\beta u^m}| \le e^{-\operatorname{Re}(\beta)u^m}|e^{-\eta\beta u^m} - 1| \le \eta|\beta|u^m e^{-\operatorname{Re}(\beta)u^m}$$

for all $|u| \le \delta n^{1/m}$. For such a δ-η pair we have

$$\mathcal{I}_1 \le \int_{|u|\le\delta n^{1/m}} \eta|\beta|u^m e^{-\operatorname{Re}(\beta)u^m}du \le \eta\int_{\mathbb{R}}|\beta|u^m e^{-\operatorname{Re}(\beta)u^m}du.$$

Since the latter integral above is finite we may consequently choose δ sufficiently small so that $\mathcal{I}_1 < \epsilon/2$. Also we know $e^{-\operatorname{Re}(\beta)u^m} \in L_1(\mathbb{R})$ and may therefore choose N sufficiently large so that $\mathcal{I}_2 < \epsilon/2$ for all $n > N$. Combining these estimates gives the desired result. □

Lemma 8. *Let $\nu : \mathbb{R} \to \mathbb{C}$ be analytic on a neighborhood of a point ξ_0 such that $|\nu(\xi_0)| = 1$. Assume that $\xi_0 \in (2;2)$ with associated constants α and β. Let $K \subseteq \mathbb{R}$ be a compact set. Then for all $\epsilon > 0$ there is a $\delta > 0$ and a natural number N such that*

$$\left|\frac{n^{1/2}}{2\pi}\int_{|\xi-\xi_0|\le\delta}\nu(\xi)^n e^{-i(xn^{1/2}+\alpha n)\xi}d\xi - e^{-i(xn^{1/2}+\alpha n)\xi_0}\nu(\xi_0)^n H_2^\beta(x)\right| < \epsilon \quad (32)$$

for all $n > N$ and for all $x \in K$.

Proof. Let $\epsilon > 0$, let $K \subseteq \mathbb{R}$ be a fixed compact set and choose $\delta > 0$ so that the estimates (17),(18) and (19) of Lemma 1 are valid. Changing variables of integration we write

$$\begin{aligned}
&\frac{n^{1/2}}{2\pi}\int_{|\xi-\xi_0|\le\delta}\nu(\xi)^n e^{-i(xn^{1/2}+\alpha n)\xi}d\xi = \\
&\qquad e^{-i(xn^{1/2}+\alpha n)\xi_0}\nu(\xi_0)^n\frac{n^{1/2}}{2\pi}\int_{|\xi|\le\delta}[\nu(\xi_0)^{-1}\nu(\xi+\xi_0)]^n e^{-i(xn^{1/2}+\alpha n)\xi}d\xi.
\end{aligned}$$

Upon setting

$$\mathcal{D} = \left| \frac{n^{1/2}}{2\pi} \int_{|\xi-\xi_0|\le\delta} \nu(\xi)^n e^{-i(xn^{1/2}+\alpha n)\xi} d\xi - e^{-i(xn^{1/2}+\alpha n)\xi_0} \nu(\xi_0)^n H_2^\beta(x) \right|,$$

we have

$$\begin{aligned} \mathcal{D} \le & \left| \frac{n^{1/2}}{2\pi} \int_{|\xi|\le Mn^{-1/2}} [\nu(\xi_0)^{-1}\nu(\xi+\xi_0)]^n e^{-i(xn^{1/2}+\alpha n)\xi} d\xi - H_2^\beta(x) \right| \\ & \qquad + n^{1/2} \left| \int_{Mn^{-1/2}<|\xi|\le\delta} [\nu(\xi_0)^{-1}\nu(\xi+\xi_0)]^n e^{-i(xn^{1/2}+\alpha n)\xi} d\xi \right|. \end{aligned}$$

where for now $0 < M < \delta n^{1/2}$ and we have used the fact that $|\nu(\xi_0)| = 1$. Continuing in this manner we have

$$\begin{aligned} \mathcal{D} \le & \left| \frac{1}{2\pi} \int_{|u|\le M} [\nu(\xi_0)^{-1}\nu(un^{-1/2}+\xi_0)]^n e^{-i(x+\alpha n^{1/2})u} du - H_2^\beta(x) \right| \\ & \qquad + n^{1/2} \left| \int_{Mn^{-1/2}<|\xi|\le\delta} [\nu(\xi_0)^{-1}\nu(\xi+\xi_0)]^n e^{-i(xn^{1/2}+\alpha n)\xi} d\xi \right| \\ \le & \left| \int_{|u|\le M} \left([\nu(\xi_0)^{-1}\nu(un^{-1/2}+\xi_0) e^{-i\alpha un^{-1/2}}]^n - e^{-\beta u^2} \right) e^{-ixu} du \right| \\ & \qquad + \left| \int_{|u|>M} e^{-ixu-\beta u^2} du \right| \\ & \qquad + n^{1/2} \left| \int_{Mn^{-1/2}}^{\delta} [\nu(\xi_0)^{-1}\nu(\xi+\xi_0)]^n e^{-i(xn^{1/2}+\alpha n)\xi} d\xi \right| \\ & \qquad + n^{1/2} \left| \int_{-\delta}^{-Mn^{-1/2}} [\nu(\xi_0)^{-1}\nu(\xi+\xi_0)]^n e^{-i(xn^{1/2}+\alpha n)\xi} d\xi \right| \\ & = \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3 + \mathcal{I}_4 \end{aligned}$$

where we have made a change of variables and used the definition of H_2^β. We now estimate the terms $\mathcal{I}_i$ for $i = 1, 2, 3, 4$. First, using Lemma 2 we may choose

$M > 0$ so that $\mathcal{I}_2 \leq \epsilon/4$ for all $x \in K$. Let us now focus on $\mathcal{I}_3$. We write

$$\begin{aligned}\mathcal{I}_3 &= n^{1/2}\left|\int_{Mn^{-1/2}}^{\delta}[\nu(\xi_0)^{-1}\nu(\xi+\xi_0)]^n e^{-i(xn^{1/2}+\alpha n)\xi}d\xi\right| \\ &= n^{1/2}\left|\int_{Mn^{-1/2}}^{\delta}[\nu(\xi_0)^{-1}\nu(\xi+\xi_0)\exp(-i\alpha+i\xi^2 p(\xi))]^n e^{i(n\xi^2 p(\xi)-\xi xn^{1/2})}d\xi\right| \\ &= n^{1/2}\left|\int_{Mn^{-1/2}}^{\delta} g(\xi)^n e^{if_n(\xi)}d\xi\right|.\end{aligned}$$

We wish to apply Lemma 5 to the above integral. Set

$$B = 4\left(1+\int_{\mathbb{R}} D|u|^{k-1}e^{-C|u|^k}du\right)$$

where $C, D \geq 0$ are the constants appearing in (18) and (19) of Lemma 1. Since $\xi^2 p(\xi)$ is a polynomial with 2 being the smallest power of its terms, we may further restrict $\delta > 0$ so that $f''(\xi) \neq 0$ and

$$c_1\xi \leq \frac{d}{d\xi}(\xi^2 p(\xi)) \leq c_2\xi$$

for all $\xi \in [Mn^{-1/2}, \delta]$ where c_1 and c_2 are non-zero real numbers of the same sign. We may consequently select $M > 0$ and a natural number N so that

$$\inf_{\xi\in[Mn^{1/2},\delta]}|f'(\xi)| > \frac{4Bn^{1/2}}{\epsilon}$$

for all x in the compact set K and for all $n > N$. Finally, an application of Lemma 5 with the above estimate and a calculation similar to that done in the proof of Lemma 6 shows

$$\mathcal{I}_3 \leq \frac{B}{\inf_{\xi\in[Mn^{1/2},\delta]}|f'(\xi)|} < \frac{\epsilon}{4}$$

for all $n > N$ and for all $x \in K$. An analogous argument gives the same estimate for $\mathcal{I}_4$.

Before treating $\mathcal{I}_1$, we fix M as the maximum M for which the above estimates will hold simultaneously. From Lemma 1 we use (17) and recall that $p(\xi)$ is continuous, $\beta = ip(0)$ and $2 < k$. We therefore observe that for any $u \in [-M, M]$

$$\lim_{n\to\infty}[\nu(\xi_0)^{-1}\nu(un^{-1/2}+\xi_0)e^{-i\alpha un^{-1/2}}]^n = e^{-iu^2 p(0)} = e^{-\beta u^2}.$$

An application of Lebesgue's dominated convergence theorem now shows that for n sufficiently large

$$\mathcal{I}_1 = \left|\int_{-M}^{M}\left([\nu(\xi_0)^{-1}\nu(un^{-1/2}+\xi_0)e^{-i\alpha un^{-1/2}}]^n - e^{-\beta u^2}\right)e^{-ixu}du\right| < \epsilon/4$$

for all $x \in \mathbb{R}$ and in particular for all $x \in K$. Finally, from the above arguments we choose $\delta > 0$ and a natural number N so that each estimate $\mathcal{I}_j < \epsilon/4$ for $j = 1, 2, 3, 4$ holds for all $n > N$ and for all $x \in K$. Putting these estimates together shows that $\mathcal{D} < \epsilon$ as desired. ☐

Lemma 9. *Let $\nu : \mathbb{R} \to \mathbb{C}$ be analytic on a neighborhood of a point ξ_0 such that $|\nu(\xi_0)| = 1$. Let $m > 2$ and assume that $\xi_0 \in (2; m)$ with associated constants α and β. Then for all $\epsilon > 0$ there is a $\delta > 0$ and a natural number N such that*

$$\left| \frac{n^{1/m}}{2\pi} \int_{|\xi - \xi_0| \le \delta} \nu(\xi)^n e^{-ix\xi} d\xi - e^{-ix\xi_0} \nu(\xi_0)^n H_m^\beta \left(\frac{x - \alpha n}{n^{1/m}} \right) \right| < \epsilon \tag{33}$$

for all $n > N$ and for all real numbers x.

The present lemma's proof will be analogous to the proof of the previous lemma in many ways. We will consequently spend less time explaining the order in which we choose our constants.

Proof. Let $\epsilon > 0$ and set

$$y_n = \frac{x - \alpha n}{n^{1/m}} \tag{34}$$

and

$$\mathcal{D} = \left| \frac{n^{1/m}}{2\pi} \int_{|\xi - \xi_0| \le \delta} \nu(\xi)^n e^{-ix\xi} d\xi - e^{-ix\xi_0} \nu(\xi_0)^n H_m^\beta \left(\frac{x - \alpha n}{n^{1/m}} \right) \right|.$$

Since $\xi_0 \in (2; m)$ we choose $\delta > 0$ so that the estimates (17), (18) and (19) of Lemma 1 are valid and the inequality

$$\left| \frac{d^2}{d\xi^2} \xi^m p(\xi) \right| \ge B^2 |\xi|^{m-2} \tag{35}$$

holds for all $-\delta \le \xi \le \delta$ where $B > 0$. Using Proposition 2 we now choose $M > 0$ such that

$$\left| \int_{|u| > M} e^{-iyu - \beta u^m} du \right| < \epsilon/4 \tag{36}$$

for all $y \in \mathbb{R}$ and

$$\frac{8}{BM^{m/2-1}} \left(1 + \int_{\mathbb{R}} D|u|^{k-1} e^{-Cu^k} du \right) < \epsilon/4 \tag{37}$$

where B was defined above and C and D are the constants appearing in (19).

As in the last lemma we write

$$\begin{aligned}\mathcal{D} &= \left|\frac{n^{1/m}}{2\pi}\int_{-\delta}^{\delta}\nu(\xi+\xi_0)^n e^{-ix(\xi+\xi_0)}d\xi - e^{-ix\xi_0}\nu(\xi_0)^n H_m^{\beta}(y)\right| \\ &\leq \left|\int_{|u|\leq M}\left([\nu(\xi_0)^{-1}\nu(un^{-1/m}+\xi_0)e^{-\alpha un^{-1/m}}]^n - e^{-\beta u^m}\right)e^{-iyu}du\right| \\ &+ \left|\int_{|u|>M} e^{-iyu-\beta u^m}du\right| + n^{1/m}\left|\int_{Mn^{-1/m}}^{\delta}[\nu(\xi_0)^{-1}\nu(\xi+\xi_0)]^n e^{-ix\xi}d\xi\right| \\ &\qquad + n^{1/m}\left|\int_{-\delta}^{-Mn^{-1/m}}[\nu(\xi_0)^{-1}\nu(\xi+\xi_0)]^n e^{-ix\xi}d\xi\right| \\ &\qquad\qquad = \mathcal{I}_1+\mathcal{I}_2+\mathcal{I}_3+\mathcal{I}_4.\end{aligned}$$

Now we estimate the terms $\mathcal{I}_j$ for $j = 1,2,3,4$. Already from (36), we have $\mathcal{I}_2 < \epsilon/4$ for all $x \in \mathbb{R}$. For $\mathcal{I}_3$ we write

$$\begin{aligned}& n^{1/m}\left|\int_{Mn^{-1/m}}^{\delta}[\nu(\xi_0)^{-1}\nu(\xi+\xi_0)]^n e^{-ix\xi}d\xi\right| \\ &= n^{1/m}\left|\int_{Mn^{-1/m}}^{\delta}[\nu(\xi_0)^{-1}\nu(\xi+\xi_0)e^{-i\alpha\xi+i\xi^m p(\xi)}]^n e^{-i(x+\alpha n)\xi-in\xi^m p(\xi)}d\xi\right| \\ &\qquad = n^{1/m}\left|\int_{Mn^{-1/m}}^{\delta} g(\xi)^n e^{if_n(\xi)}d\xi\right|\end{aligned}$$

where

$$g(\xi) = [\nu(\xi_0)^{-1}\nu(\xi+\xi_0)e^{-i\alpha\xi+i\xi^m p(\xi)}]$$

and

$$f_n(\xi) = -[(x+\alpha n)\xi - in\xi^m p(\xi)].$$

With the aim of applying Lemma 5, we use (35) and observe that on the interval $[Mn^{-1/m},\delta]$

$$\inf|f_n''(\xi)| \geq \inf nB^2|\xi|^{m-2} \geq nB^2|Mn^{-1/m}|^{m-2} = (n^{1/m}BM^{m/2-1})^2 > 0.$$

The application of the lemma is therefore justified and we may use (18) and (19) to see that

$$\begin{aligned}\mathcal{I}_3 &\leq \frac{8n^{1/m}}{\sqrt{(n^{1/m}BM^{m/2-1})^2}}(||g^n||_\infty + ||ng'g^{n-1}||_1) \\ &\qquad \leq \frac{8}{CM^{m/2-1}}\left(1+\int_{\mathbb{R}} D|u|^{k-1}e^{-Cu^k}du\right) < \epsilon/4\end{aligned}$$

for all $x \in \mathbb{R}$. A similar calculation gives the same estimate for $\mathcal{I}_4$.

Finally we treat $\mathcal{I}_1$. Exactly as we did in the previous lemma we use (17) and observe that for any $u \in [-M, M]$

$$\lim_{n\to\infty}[\nu(\xi_0)^{-1}\nu(un^{-1/m}+\xi_0)e^{-\alpha u n^{-1/m}}]^n = e^{-\beta u^m}.$$

An application of Lebesgue's dominated convergence theorem gives a natural number N for which $\mathcal{I}_1 \leq \epsilon/4$ for all $n > N$ and for all $x \in \mathbb{R}$. Combining our estimates finishes the proof. $\square$

We are now in a position to prove the local limit theorems discussed in the introduction. Let $\phi : \mathbb{Z} \to \mathbb{C}$ have admissible support and let $\hat{\phi}$ be its Fourier transform. Define $\tilde{\phi} : \mathbb{N} \times \mathbb{R} \to \mathbb{Z}$ by

$$\tilde{\phi}(n,x) = \frac{1}{2\pi}\int_{-\pi}^{\pi}\hat{\phi}(\xi)^n e^{-ix\xi}d\xi. \tag{38}$$

It is easy to see that $\tilde{\phi}(n,\cdot)$ is an analytic function. By Properties of the Fourier transform

$$\tilde{\phi}(n,x) = \phi^{(n)}(x) \tag{39}$$

for all $x \in \mathbb{Z}$, i.e. $\tilde{\phi}(n,\cdot)$ is a smooth extension of $\phi^{(n)}$.

In the remainder of this section we state and prove two theorems after which their results are combined to prove Theorem 2. The proof of the first, Theorem 5, is the most technical; it corresponds to the case where $\phi^{(n)}$ is approximated by the heat kernel evaluated at purely imaginary time. Theorem 6 isolates the second conclusion, (10), of Theorem 3.

Theorem 5. *Let $\phi : \mathbb{Z} \to \mathbb{C}$ have admissible support. Suppose that Fourier transform of ϕ satisfies* $\sup|\hat{\phi}(\xi)| = 1$ *and that for every $\rho \in (-\pi,\pi]$ such that $|\hat{\phi}(\rho)| = 1$ we have*

$$\beta(\rho) = \frac{1}{2}\frac{d^2}{d\xi^2}\log(\hat{\phi}(\xi))\Big|_{\rho} \neq 0 \tag{40}$$

and for at least one such ρ, $\beta(\rho)$ is purely imaginary. Then there is a finite collection of real numbers $\alpha_1, \alpha_2, \dots, \alpha_Q$ such that for every α_q there is a finite collection of points $\xi_1, \xi_2, \dots, \xi_{R_q}$ and complex numbers $\beta_1, \beta_2, \dots, \beta_{R_q}$ such that

$$\phi^{(n)}(\lfloor xn^{1/2}+\alpha_q n\rfloor) = \sum_{l=1}^{R_q} n^{-1/2}e^{-i(\lfloor xn^{1/2}+\alpha_q n\rfloor)\xi_l}\hat{\phi}(\xi_l)^n H_2^{\beta_l}(x) + o(n^{-1/2}) \tag{41}$$

uniformly on any compact set.

Proof. We let $\epsilon > 0$ and $K \subseteq \mathbb{R}$ be a compact set. Let

$$\Omega = \{\rho \in (-\pi,\pi] : |\hat{\phi}(\rho)| = 1\}.$$

It follows from Proposition 1 and (40) that Ω is a finite set and for each $\rho \in \Omega$, $\rho \in (1;2)$ or $\rho \in (2;2)$. This is true precisely because $\beta(\rho)$ is the coefficient of the ξ^2 term in the Taylor series expansion for η, and by our assumption is non-zero. We take the set real numbers $\alpha_1, \alpha_2, \ldots, \alpha_Q$ to be the associated drift constants for the subset of points $\rho \in \Omega$ for which $\beta(\rho)$ is purely imaginary. Equivalently $\alpha_1, \alpha_2, \ldots, \alpha_Q$ are the associated drift constants for the points $\rho \in \Omega$ which are of type 2 for $\hat{\phi}$. By our hypothesis, at least one such α_q exists.

Let α_q be a member of the above collection. We will index the set $\Omega = \{\xi_1, \xi_2, \ldots, \xi_P\}$ as follows:

- For $1 \leq l \leq \widetilde{R}_q$, $\xi_l \in (2;2)$ with associated constants, $k_l, \alpha_l = \alpha_q, \gamma_l$ and real polynomial $p_l(\xi)$ where we have set $\beta_l = ip_l(0)$
- For $\widetilde{R}_q < l \leq R_q$, $\xi_l \in (1;2)$, with associated constants, $\alpha_l = \alpha_q$ and β_q
- For $R_q < l \leq \widetilde{P}$, $\xi_l \in (2;2)$ with associated constants, $k_l, \alpha_l \neq \alpha_q, \gamma_l$ and real polynomial $p_l(\xi)$ where we have set $\beta_l = ip_l(0)$
- For $\widetilde{P} < l \leq P$, $\xi_l \in (1;2)$ with associated constants, $\alpha_l \neq \alpha_q$ and β_l

Here $\widetilde{R}_q \leq R_q \leq \widetilde{P} \leq P$. We remark that only one of the above subcollections of $\xi_1, \xi_2, \ldots, \xi_P$ is necessarily non-empty, for it might be the case that $1 = R_q = P$ and in this case the result (41) simplifies considerably. In fact, this is precisely the situation exemplified in the introduction.

We divide the interval $(-\pi, \pi]$ into subintervals. For $l = 1, 2, \ldots, P$ let $I_l = [\xi_l - \delta_l, \xi_l + \delta_l]$ for values $\delta_l > 0$ to be chosen later and let $J = (-\pi, \pi]/\cup I_l$. Write

$$\begin{aligned}
n^{1/2}\tilde{\phi}(n, xn^{1/2} + \alpha_q n) &= \frac{n^{1/2}}{2\pi}\int_{-\pi}^{\pi} \hat{\phi}(\xi)^n e^{-i(xn^{1/2}+\alpha_q n)\xi} d\xi \\
&= \sum_{l=1}^{P} \frac{n^{1/2}}{2\pi}\int_{I_l} \hat{\phi}(\xi)^n e^{-i(xn^{1/2}+\alpha_q n)\xi} d\xi \\
&+ \frac{n^{1/2}}{2\pi}\int_{J} \hat{\phi}(\xi)^n e^{-i(xn^{1/2}+\alpha_q n)\xi} d\xi \\
&= \sum_{l=1}^{P} \mathcal{I}_l + \mathcal{E}. \qquad (42)
\end{aligned}$$

We treat the integrals $\mathcal{I}_l$ in the four cases separately. First we consider $\mathcal{I}_1$

where $l = R_q + 1, R_q + 2, \ldots, \widetilde{P}$. Here

$$\begin{aligned}
|\mathcal{I}_l| &= \left| \frac{n^{1/2}}{2\pi} \int_{I_l} \hat{\phi}(\xi)^n e^{-i(xn^{1/2}+\alpha_q n)\xi} d\xi \right| \\
&= \left| \frac{n^{1/2} e^{-i(xn^{1/2}-\alpha_q n)\xi_l} \hat{\phi}(\xi_l)^n}{2\pi} \int_{|\xi|\le\delta} \left[\hat{\phi}(\xi_l)^{-1} \hat{\phi}(\xi_l + \xi) \right]^n e^{-i(xn^{1/2}+\alpha_q n)\xi} d\xi \right| \\
&\le n^{1/2} \left| \int_{|\xi|\le\delta} \left[\hat{\phi}^{-1}(\xi_l) \hat{\phi}(\xi_l + \xi) e^{-i\alpha_l \xi + i\xi^2 p_l(\xi)} \right]^n e^{-in[(xn^{-1/2}+\alpha_q-\alpha_l)\xi+\xi^2 p_l(\xi)]} d\xi \right|.
\end{aligned}$$

Set

$$g_l(\xi) = [\hat{\phi}^{-1}(\xi_l)\hat{\phi}(\xi_l + \xi)e^{-i\alpha_l\xi + i\xi^2 p_l(\xi)}]$$

and

$$f_{n,l}(\xi) = -n[(xn^{-1/2} + \alpha_q - \alpha_l)\xi + \xi^2 p_l(\xi)]$$

We may therefore write

$$|\mathcal{I}_q| \le n^{1/2} \left| \int_{|\xi|\le\delta} g_l(\xi)^n e^{if_{n,l}(\xi)} d\xi \right|.$$

To estimate the above integral, we choose $\delta_l > 0$ so that the on the interval $[-\delta_l, \delta_l]$, our function $g_l(\xi)$ satisfies (18) and (19) for some $C_l, D_l > 0$,

$$f''_{n,l}(\xi) = -n\frac{d^2}{d\xi^2}\xi^2 p_l(\xi) \neq 0$$

and

$$B_l \le |\alpha_l - \alpha_q - \frac{d}{d\xi}\xi^2 p_l(\xi)| \tag{43}$$

for some $B_l > 0$. For the first property our choice of δ_l was made using Lemma 1 and the assumption that $\xi_l \in (2; 2)$. For the second two properties we used that fact that $\xi^2 p_l(\xi)$ is a polynomial with 2 being the smallest power of its terms and $\alpha_l \neq \alpha_q$. We may therefore apply Lemma 5. This gives

$$\begin{aligned}
|\mathcal{I}_l| &\le \frac{8n^{1/2}}{\inf_\xi |f'_{n,l}(\xi)|}(||g_l||_\infty + ||ng'_l g_l^{n-1}||) \\
&\le \frac{8}{\inf_\xi |(x - n^{1/2}(\alpha_l - \alpha_q - \frac{d}{d\xi}\xi^2 p_l(\xi))|} \left(1 + \int_{\mathbb{R}} D_l |\xi|^{k_l} e^{-C_l \xi^{k_l}} d\xi \right) \\
&\le \frac{M_l}{\inf_\xi |(x - n^{1/2}(\alpha_l - \alpha_q - \frac{d}{d\xi}\xi^2 p_l(\xi))|}
\end{aligned}$$

for some $M_l > 0$ and where the above infima are taken over the interval $[-\delta_l, \delta_l]$. Using the estimate (43) and recalling that x lives inside the compact set K we may choose a natural number N_l so that

$$\inf_{\xi} |(x - n^{1/2}(\alpha_l - \alpha_q - \frac{d}{d\xi}\xi^2 p_l(\xi))| > \frac{M_l(P+1)}{\epsilon}$$

for all $n > N_l$ and for all $x \in K$. Consequently,

$$|\mathcal{I}_l| \leq \frac{M_l}{M_l(P+1)/\epsilon} = \epsilon/(P+1) \tag{44}$$

for all $n > N_l$ and for all $x \in K$.

We now consider the integrals $\mathcal{I}_l$ where $l = \widetilde{P}+1, \widetilde{P}+2, \ldots, P$. Appealing to Lemma 7 we may choose $\delta_l > 0$ and a natural number N_l so that

$$\begin{aligned} |\mathcal{I}_l| &\leq \epsilon/2(P+1) + |e^{-i(xn^{1/2}+\alpha_q n)\xi_l}\hat{\phi}(\xi_l)^n H_2^{\beta_l}(x + (\alpha_q - \alpha_l)n^{1/2})| \\ &\leq \epsilon/2(P+1) + |H_2^{\beta_l}(x + (\alpha_q - \alpha_l)n^{1/2})| \end{aligned}$$

for all $n > N_l$ and for all $x \in \mathbb{R}$. However, as we remarked earlier $H_2^{\beta_l}$ is the heat kernel evaluated at complex time β_l. Since $\mathrm{Re}(\beta_l) > 0$ in this case and $\alpha_q \neq \alpha_l$ we may increase our natural number N_l to ensure that

$$|H_2^{\beta_l}(x + (\alpha_q - \alpha_l)n^{1/2})| \leq \epsilon/2(P+1)$$

for any $n > N_l$ and for all x in the compact set K. These estimates together give

$$|\mathcal{I}_l| \leq \epsilon/2(P+1) + \epsilon/2(P+1) = \epsilon/(P+1) \tag{45}$$

for all $n > N_l$ and for all $x \in K$.

In the remaining estimates of $\mathcal{I}_l$ for $l = 1, 2, \ldots, R_q$ we recall that $\alpha_q = \alpha_l$. For the integrals $\mathcal{I}_l$ for $l = 1, 2, \ldots, \widetilde{R}_q$ we appeal to Lemma 8. From this we choose $\delta_l > 0$ and a natural number N_l such that

$$|\mathcal{I}_l - e^{-i(xn^{1/2}+\alpha_q n)\xi_l}\hat{\phi}(\xi_l)^n H_2^{\beta_l}(x)| \leq \epsilon/(P+1) \tag{46}$$

for all $n > N_l$ and for all $x \in K$.

Finally we treat the integrals $\mathcal{I}_l$ for $l = \widetilde{R}_q + 1, \widetilde{R}_q + 2, \ldots, R_q$. Here we appeal to Lemma 7 and chose $\delta_l > 0$ and N_l, a natural number, such that

$$|\mathcal{I}_l - e^{-i(xn^{1/2}+\alpha_q n)\xi_l}\hat{\phi}(\xi_l)^n H_2^{\beta_l}(x)| \leq \epsilon/(P+1) \tag{47}$$

for all $n > N_l$ and for all $x \in \mathbb{R}$. In particular we have this estimate uniform for all $x \in K$.

After fixing our collection of δ_l's in the above arguments, the set J becomes fixed. We therefore set $s = \sup_{\xi \in J} |\hat{\phi}(\xi)| < 1$ and note that $|\mathcal{E}| \leq n^{1/2}s^n$. Thus we may choose a natural number N_0 such that $|\mathcal{E}| < \epsilon/(P+1)$ for all $n > N_0$ and for all $x \in K$.

At last, we choose N to be the maximum of N_l for $l = 0, 1, \ldots, P$. Combining the estimates (42), (44), (45), (46) and (47) yields

$$\begin{aligned}\left| n^{1/2}\tilde{\phi}(n, xn^{1/2} + \alpha_q n) - \sum_{l=1}^{P} e^{-i(xn^{1/2}+\alpha_q n)\xi_q} \hat{\phi}(\xi_q)^n H_2^{\beta_q}(x) \right| \\ \leq \sum_{l=1}^{R_q} \left| \mathcal{I}_l - e^{-i(xn^{1/2}+\alpha_q n)\xi_q} \hat{\phi}(\xi_l)^n H_2^{\beta_l}(x) \right| + \sum_{R_q < l \leq P} |\mathcal{I}_l| + \mathcal{E} \\ < \frac{(P+1)\epsilon}{P+1} = \epsilon\end{aligned}$$

for any $n > N$ and for all $x \in K$. We have shown that

$$\tilde{\phi}(n, xn^{1/2} + \alpha_q n) = \sum_{l=1}^{R_q} n^{-1/2} e^{-i(xn^{1/2}+\alpha_q n)\xi_l} \hat{\phi}(\xi_l)^n H_2^{\beta_l}(x) + o(n^{-1/2}) \tag{48}$$

uniformly for x in any compact set K.

To complete the proof of the theorem we need to replace the argument $xn^{1/2} + \alpha_q n$ by an integer in (48); this is precisely where the floor function comes in. Let K be compact, set

$$y(x, n) = \frac{\lfloor \alpha_q n + xn^{1/2} \rfloor - \alpha_q n}{n^{1/2}}$$

and observe that $|x - y(n, x)| \leq n^{-1/2}$. Let $F \supseteq K$ be any compact set for which $y(x, n) \in F$ for all $x \in K$ and all natural numbers n. By Proposition 4, each function $H_2^{\beta_l}$ is uniformly continuous on F and therefore for any $x \in K$ we have

$$\begin{aligned}\sum_{l=1}^{R_q} n^{-1/2} e^{-i(\lfloor xn^{1/2}+\alpha_q n \rfloor)\xi_l} \hat{\phi}(\xi_l)^n H_2^{\beta_l}(y(x, n)) \\ = \sum_{l=1}^{R_q} n^{-1/2} e^{-i(\lfloor xn^{1/2}+\alpha_q n \rfloor)\xi_l} \hat{\phi}(\xi_l)^n H_2^{\beta_l}(x) + o(n^{-1/2}).\end{aligned} \tag{49}$$

The result now follows from (48), (49) and the observation that

$$\phi^{(n)}(\lfloor xn^{1/2} + \alpha_q n \rfloor) = \tilde{\phi}(n, \lfloor xn^{1/2} + \alpha_q n \rfloor).$$

□

Theorem 6. *Let $\phi : \mathbb{Z} \to \mathbb{C}$ have admissible support. Suppose that the Fourier transform of ϕ satisfies* $\sup |\hat{\phi}(\xi)| = 1$ *and for every $\rho \in (-\pi, \pi]$, such that* $|\hat{\phi}(\rho)| = 1$

$$\frac{d^2}{d\xi^2} \log(\hat{\phi}(\xi)) \Big|_{\rho} \neq i\tau \tag{50}$$

for any non-zero real number τ. Then there exist a natural number $m \geq 2$, a finite number of points $\xi_1, \xi_2, \ldots, \xi_Q$, real numbers $\alpha_1, \alpha_2, \ldots, \alpha_Q$ and complex numbers $\beta_1, \beta_2, \ldots, \beta_Q$ such that

$$\phi^{(n)}(x) = \sum_{q=1}^{Q} n^{-1/m} e^{-ix\xi_q} \hat{\phi}(\xi_q)^n H_m^{\beta_q}\left(\frac{x - \alpha_q n}{n^{1/m}}\right) + o(n^{-1/m}) \tag{51}$$

uniformly in $\mathbb{Z}$.

Proof. As in the proof of the previous theorem we first appeal to Proposition 1. The proposition gives a finite number of points $\xi_1, \xi_2, \ldots, \xi_R \in (-\pi, \pi]$ at which $|\hat{\phi}(\xi)| = 1$. Moreover $\xi_q \in (1; m_q)$ or $(2; m_q)$ for each $q = 1, 2, \ldots, R$. Let $m = \max_q\{m_q\}$. After a possible reindex, we will assume the set of points $\xi_1, \xi_2, \ldots, \xi_R$ is organized as follows:

- For $1 \leq q \leq Q$, $\xi_q \in (1; m)$ or $\xi_q \in (2; m)$ with associated constants α_q and β_q
- If $Q < R$ and $Q < q \leq R$, ξ_q is a point of order m_q where $m_q < m$

Comparing our hypothesis (50) to the Taylor series for η in definition 1, we conclude that either $m > 2$ or, in the case that $m = 2$ we must have $\xi_q \in (1; 2)$ for each q. Consequently to each point ξ_q of order m we may apply either Lemma 7 or Lemma 9.

Let $\epsilon > 0$. As in the proof of the previous theorem we divide the interval $(-\pi, \pi]$ into subintervals. For $q = 1, 2, \ldots, R$ let $I_q = [\xi_q - \delta_q, \xi_q + \delta_q]$ for values of δ_q to be chosen later and let $J = (-\pi, \pi] \setminus \cup I_q$. Write

$$\begin{aligned} n^{1/m}\tilde{\phi}(n, x) &= \frac{n^{1/m}}{2\pi}\int_{-\pi}^{\pi} \hat{\phi}(\xi)^n e^{-ix\xi} d\xi \\ &= \sum_{q=1}^{R} \frac{n^{1/m}}{2\pi}\int_{I_q} \hat{\phi}(\xi)^n e^{-ix\xi} d\xi \\ &+ \frac{n^{1/m}}{2\pi}\int_{J} \hat{\phi}(\xi)^n e^{-ix\xi} d\xi. \end{aligned}$$

Therefore,

$$\begin{aligned} &\left| n^{1/m}\tilde{\phi}(n, x) - \sum_{q=1}^{Q} e^{-ix\xi_q}\hat{\phi}(\xi_q)^n H_m^{\beta_q}\left(\frac{x - \alpha_q n}{n^{1/m}}\right)\right| \\ &\leq \sum_{q=1}^{Q}\left|\frac{n^{1/m}}{2\pi}\int_{I_q}\hat{\phi}(\xi)^n e^{-ix\xi} d\xi - e^{-ix\xi_q}\hat{\phi}(\xi_q)^n H_m^{\beta_q}\left(\frac{x - \alpha_q n}{n^{1/m}}\right)\right| \\ &+ \sum_{q=Q+1}^{R} n^{1/m}\left|\frac{1}{2\pi}\int_{I_q}\hat{\phi}(\xi)^n e^{-ix\xi} d\xi\right| + \left|\frac{n^{1/m}}{2\pi}\int_{J}\hat{\phi}(\xi)^n e^{-ix\xi} d\xi\right|. \end{aligned} \tag{52}$$

For $q = 1, 2, \dots, Q$ we apply either Lemma 7 or Lemma 9. We may therefore choose a natural number N_q and fix $\delta_q > 0$ so that

$$\left| \frac{n^{1/2}}{2\pi} \int_{I_q} \hat{\phi}(\xi)^n e^{-ix\xi} d\xi - e^{-ix\xi_q} \hat{\phi}(\xi_q)^n H_m^{\beta_q} \left(\frac{x - \alpha_q n}{n^{1/m}} \right) \right| < \frac{\epsilon}{(R+1)} \tag{53}$$

for all $n > N_q$ and for all $x \in \mathbb{R}$.

In the case that $q = Q+1, Q+2, \dots, R$ we appeal to Lemma 6 and choose $\delta_q > 0$ and a natural number N_q such that

$$\left| \frac{1}{2\pi} \int_{I_q} \hat{\phi}(\xi)^n e^{-ix\xi} d\xi \right| \le \frac{C_q}{n^{1/m_q}}$$

for some $C_q > 0$ and for all $n > N_q$ and $x \in \mathbb{R}$. Using the fact that $m > m_q$ we may adjust the value of N_q so that

$$n^{1/m} \left| \frac{1}{2\pi} \int_{I_q} \hat{\phi}(\xi)^n e^{-ix\xi} d\xi \right| \le \frac{C_q}{n^{1/m_q - 1/m}} < \frac{\epsilon}{(R+1)} \tag{54}$$

for all $n > N_q$ and for all $x \in \mathbb{R}$.

Finally, as in the proof of the last theorem, we set $s = \inf_J |\hat{\phi}| < 1$ and observe that the last term in (52) is bounded by $n^{1/m} s^n$. We therefore select a natural number N_0 such that

$$\left| \frac{n^{1/2}}{2\pi} \int_J \hat{\phi}(\xi)^n e^{-ix\xi} d\xi \right| \le n^{1/m} s^n < \frac{\epsilon}{(R+1)} \tag{55}$$

for all $n > N_0$ and for all $x \in \mathbb{R}$.

Let us choose N to be the maximum N_q for $q = 0, 1, \dots, R$. Upon combining the estimates (53), (54), (55) and (52) we have

$$\left| n^{1/m} \tilde{\phi}(n, x) - \sum_{q=1}^{Q} e^{-ix\xi_q} \hat{\phi}(\xi_q)^n H_m^{\beta_q} \left(\frac{x - \alpha_q n}{n^{1/m}} \right) \right| < \epsilon \tag{56}$$

for all $n > N$ and for all $x \in \mathbb{R}$. In particular, it holds for all $x \in \mathbb{Z}$ and in which case $\tilde{\phi}(n, x) = \phi^{(n)}(x)$. This is our desired result. □

Proof of Theorem 2. Let K be a compact set. Assuming that ϕ satisfies the hypotheses of the theorem, we consider its Fourier transform $\hat{\phi}$. We define

$$\beta(\rho) = \frac{1}{2} \frac{d^2}{d\xi^2} \log(\hat{\phi}(\xi)) \Big|_{\rho} \tag{57}$$

for $\rho \in (-\pi, \pi]$. There are two possibilities for $\beta(\rho)$, either $\beta(\rho)$ satisfies the hypotheses of Theorem 5 or $\beta(\rho)$ satisfies the hypotheses of Theorem 6. A moments thought shows that the hypotheses are indeed mutually exclusive and collectively exhaustive. If the case at hand is the former there is nothing to prove for $m = 2$ and the desired result is precisely the conclusion of Theorem 5. We will therefore address the latter case.

From Theorem 6 take $\alpha_1, \alpha_2, \dots, \alpha_Q$ and $m \geq 2$. Let α_q be a member of the preceding collection. We reindex the collections $\beta_1, \beta_2, \dots, \beta_Q$ and $\xi_1, \xi_2, \dots, \xi_Q$, taken from Theorem 6, as follows. Let $\beta_1, \beta_2, \dots, \beta_{R_q}$ and $\xi_1, \xi_2, \dots, \xi_{R_q}$ be such that the corresponding drift constants, $\alpha_1, \alpha_2, \dots, \alpha_{R_q}$, satisfy $\alpha_l = \alpha_q$ for $l = 1, 2, \dots, R_q$. Further we let $\beta_{R_q+1}, \beta_{R_q+2}, \dots, \beta_Q$ and $\xi_{R_q+1}, \xi_{R_q+2}, \dots, \xi_Q$ be such that the corresponding drift constant satisfy $\alpha_l \neq \alpha_q$ for $l = R_q+1, R_q+2, \dots, Q$. Observe that (56) is uniform in $\mathbb{R}$ and we may therefore write

$$\begin{aligned}
&\tilde{\phi}(n, \alpha_q n + x n^{1/m}) \\
&\qquad = \sum_{l=1}^{Q} n^{-1/m} e^{-i(\alpha_q n + x n^{-1/m})\xi_l} \hat{\phi}(\xi_l)^n H_m^{\beta_l}\left(\frac{(\alpha_q - \alpha_l)n + x n^{1/m}}{n^{1/m}}\right) + o(n^{-1/m}) \\
&\qquad\qquad = \sum_{l=1}^{R_q} n^{-1/m} e^{-i(\alpha_q n + x n^{-1/m})\xi_l} \hat{\phi}(\xi_l)^n H_m^{\beta_l}(x) \\
&+ \sum_{l=R_q+1}^{Q} n^{-1/m} e^{-i(\alpha_q n + x n^{-1/m})\xi_l} \hat{\phi}(\xi_l)^n H_m^{\beta_l}((\alpha_q - \alpha_l) n^{1-1/m} + x) + o(n^{-1/m}). \\
&\quad = \sum_{l=1}^{R_q} n^{-1/m} e^{-i(\alpha_q n + x n^{-1/m})\xi_l} \hat{\phi}(\xi_l)^n H_m^{\beta_l}(x) + \sum_{l=R_q+1}^{Q} S_l(n, x) + o(n^{-1/m}).
\end{aligned}$$

Upon requiring $x \in K$ we consider the summands $S_l(n, x)$ for $l = R_q + 1, R_q + 2, \dots, Q$. In the case that $\mathrm{Re}(\beta_l) > 0$ we have

$$\begin{aligned}
|S_l(n, x)| &= |n^{-1/m} e^{-i(\alpha_q n + x n^{-1/m})\xi_l} \hat{\phi}(\xi_l)^n H_m^{\beta_l}((\alpha_q - \alpha_l) n^{1-1/m} + x)| \\
&= n^{-1/m} |H_m^{\beta_l}((\alpha_q - \alpha_l) n^{1-1/m} + x))| \\
&\leq n^{-1/m} C_l \exp(-B_l((\alpha_q - \alpha_l) n^{1-1/m} + x)^{m/(m-1)}) = o(n^{-1/m}).
\end{aligned}$$

If it is the case that $\mathrm{Re}(\beta_l) = 0$ we first observe that $m > 2$; this follows from the assumption that $\beta(\rho)$ satisfies the hypotheses of Theorem 6 but not Theorem 5. Appealing to Proposition 3 we conclude that

$$\begin{aligned}
&|S_l(n, x)| \\
&\quad \leq n^{-1/m} \left(\frac{A}{|(\alpha_q - \alpha_l) n^{1-1/m} + x)|^{\frac{m-2}{2(m-1)}}} + \frac{B}{|(\alpha_q - \alpha_l) n^{1-1/m} + x)|} \right) \\
&\qquad = o(n^{-1/m}).
\end{aligned}$$

Combining the above estimates shows that for all $x \in K$

$$\tilde{\phi}(n, \alpha_q n + x n^{1/m}) = \sum_{l=1}^{R_q} n^{-1/m} e^{-i(\alpha_q n + x n^{-1/m})\xi_l} \hat{\phi}(\xi_l)^n H_m^{\beta_l}(x) + o(n^{-1/m}).$$

To complete the proof, it remains to replace the argument $\alpha_q n + xn^{1/m}$ by the integer $\lfloor \alpha_q n + xn^{1/m} \rfloor$ in the equation above. This can be done easily by making an argument analogous to that given in the last paragraph of the proof to Theorem 5. From this, the desired result follows without trouble. $\square$

6 The lower bound of $||\phi^{(n)}||_\infty$

In this section we complete the proof of Theorem 1.

Lemma 10. *Let $\xi_1, \xi_2, \cdots, \xi_R \in (-\pi, \pi]$ be distinct let $B > 0$ and define*

$$V = \begin{pmatrix} 1 & 1 & \dots & 1 \\ e^{-i\xi_1} & e^{-i\xi_2} & \dots & e^{-i\xi_R} \\ e^{-i2\xi_1} & e^{-i2\xi_2} & \dots & e^{-i2\xi_R} \\ \vdots & \vdots & \ddots & \vdots \\ e^{-i(R-1)\xi_1} & e^{-i(R-1)\xi_2} & \dots & e^{-i(R-1)\xi_R} \end{pmatrix}. \tag{58}$$

Then there is a number $C_0 > 0$ such that for any $\rho, \sigma \in \mathbb{C}^R$ with $||\rho|| > B$ and $\sigma = V\rho$ we have $|\sigma_j| > 3C_0$ for some $j = 1, 2, \dots, R$. Here $||\cdot||$ denotes the usual norm on $\mathbb{C}^R$.

Proof. The matrix V in (58) is known as Vandermonde's matrix. It is a routine exercise in linear algebra to show that

$$\det(V) = \prod_{1 \le l < k \le R} (e^{-i\xi_k} - e^{-i\xi_l}).$$

Noting that $e^{-i\xi_1}, e^{-i\xi_1}, \dots, e^{-i\xi_R}$ are all distinct we conclude that V is invertible. The proof now follows immediately from the estimate

$$||\rho|| \le ||V^{-1}||||\sigma||.$$

$\square$

Proof of Theorem 1. Let $\phi : \mathbb{Z} \to \mathbb{C}$ have admissible support. As Theorem 4 gave the upper bound,

$$A^{-n}||\phi^{(n)}||_\infty \le C' n^{1/m}$$

for some $C' > 0$, our job is establish the lower bound

$$Cn^{1/m} \le A^{-n}||\phi^{(n)}||_\infty$$

for some $C > 0$. We will do this using our local limit theorems.

As we noted in the proof of Theorem 4, it suffices to assume that $A = \sup_\xi |\hat{\phi}(\xi)| = 1$. We let $m \ge 2$ be chosen as it was in the proof Theorem 2 or Theorem 4, a moments thought shows that these arguments yield the same m. By Theorem 2 we select $\alpha = \alpha_1$, complex numbers $\beta_1, \beta_2, \dots, \beta_R$ and distinct points $\xi_1, \xi_2, \dots, \xi_R \in (-\pi, \pi]$ for which the conclusion of the theorem holds.

Appealing to Proposition 4, we know that each function $H_m^{\beta_l}$ is non-zero and continuous. Thus for $H_m^{\beta_1}$ there is $B > 0$ and some interval $I = [a, b]$ for which $|H_m^{\beta_l}(x)| \geq B$ for all $x \in I$.

Define V by (58) and let $C_0 > 0$ as guaranteed by Lemma 10. Set

$$f(n, x) = \sum_{l=1}^{R} e^{-i(\alpha n + xn^{1/m})\xi_l} \hat{\phi}(\xi_l)^n H_m^{\beta_l}(x) \tag{59}$$

and

$$\sigma_k(n, x) = \sum_{l=1}^{R} e^{-ik\xi} e^{-i(\alpha n + xn^{1/m})\xi_l} \hat{\phi}(\xi_l)^n H_m^{\beta_l}(x) \tag{60}$$

for $k = 0, 1, \ldots, R-1$. Since each function $H_m^{\beta_l}$ is continuous on $\mathbb{R}$ it is uniformly continuous on $[a-R, b+R] \supseteq I$. Consequently we may choose a natural number N for which

$$|f(n, x + kn^{-1/m}) - \sigma_k(n, x)| < C_0 \tag{61}$$

for all $n \geq N$, $k = 0, 1, \ldots, R-1$ and $x \in I$. By possibly further increasing N we can also guarantee that for any $n \geq N$ there is $x_0 \in I$ such that $\alpha n + x_0 n^{1/m}$ is an integer and for which $x_0 + kn^{-1/m} \in I$ for all $k = 0, 1, \ldots, R-1$. We observe that for any such k, $(\alpha n + (x_0 + kn^{-1/m})n^{1/m})$ is also an integer.

Now for any $n \geq N$ let $x_0 \in I$ be as guaranteed in the previous paragraph. Observe that

$$\begin{array}{ccc} \begin{pmatrix} \sigma_0(n, x_0) \\ \sigma_1(n, x_0) \\ \vdots \\ \sigma_{(R-1)}(n, x_0) \end{pmatrix} & = & \begin{pmatrix} 1 & 1 & \cdots & 1 \\ e^{-i\xi_1} & e^{-i\xi_2} & \cdots & e^{-i\xi_R} \\ e^{-i2\xi_1} & e^{-i2\xi_2} & \cdots & e^{-i2\xi_R} \\ \vdots & \vdots & \ddots & \vdots \\ e^{-i(R-1)\xi_1} & e^{-i(R-1)\xi_2} & \cdots & e^{-i(R-1)\xi_n} \end{pmatrix} \begin{pmatrix} \rho_1(n, x_0) \\ \rho_2(n, x_0) \\ \vdots \\ \rho_R(n, x_0) \end{pmatrix} \\ \sigma(n, x_0) & = & V \qquad\qquad\qquad \rho(n, x_0) \end{array}$$

where

$$\rho_l(x_0, n) = e^{-i(\alpha n + x_0 n^{1/m})\xi_l} \hat{\phi}(\xi_l)^n H_m^{\beta_l}(x_0)$$

for $l = 1, 2, \ldots, R$. Because $x_0 \in I$, $|\rho_1(x_0, n)| = |H_m^{\beta_1}(x_0)| > B$ and therefore

$$||\rho(n, x_0)|| > B.$$

Appealing to Lemma 10, there is some $k \in \{0, 1, 2, \ldots, R-1\}$ such that $|\sigma_k(n, x_0)| > 3C_0$ so by (61), $|f(n, x_0 + kn^{-1/m}| > 2C_0$.

We have shown that there is a natural number N, a closed interval I and a constant $C_0 > 0$ such that for any $n \geq N$

$$\sup \left| \sum_{l=1}^{R} e^{-i(\alpha n + xn^{1/m})\xi_l} \hat{\phi}(\xi_l)^n H_m^{\beta_l}(x) \right| > 2C_0 \tag{62}$$

where the above supremum is taken over the set

$$\{x : x \in I \text{ and } (\alpha n + x n^{-1/m}) \in \mathbb{Z}\}.$$

Combining (7) and (62) we conclude that

$$\sup_{x\in\mathbb{Z}} |\phi^{(n)}(x)| \geq C n^{-1/m} \tag{63}$$

for all $n > N$. The result now follows from the observation that $\phi^{(n)} \neq 0$ for all $n \leq N$ and so, by possibly adjusting C, (63) must hold for all n. □

7 Concentration of mass

In this section we complete the proof of Theorem 3. Recall that the theorem has two conclusions, the second of which is the subject of Theorem 6 and was already shown in the previous section. What remains to be shown is the first conclusion.

Proof of Theorem 3. We assume that ϕ satisfies the hypotheses of the theorem. By Theorem 6 we have an integer $m \geq 2$, and real numbers $\alpha_1, \alpha_2, \dots, \alpha_Q$ such that

$$\phi^{(n)}(x) = \sum_{q=1}^{Q} n^{-1/m} e^{-ix\xi_q} \hat{\phi}(\xi_q)^n H_m^{\beta_q}\left(\frac{x-\alpha_q n}{n^{1/m}}\right) + o(n^{-1/m}) \tag{64}$$

where the limit is uniform for $x \in \mathbb{Z}$ and $\beta_1, \beta_2, \dots, \beta_q$ and $\xi_1, \xi_2, \dots, \xi_Q$ are associated with $\alpha_1, \alpha_2, \dots, \alpha_Q$.

Using Theorem 1 we choose $C > 0$ for which the estimate (3) holds. Considering all possibilities of β_q and m above, we can choose $M > 0$ such that

$$|H_m^{\beta_q}(y)| < C/(2Q)$$

for all $|y| > M$ and for all $q = 1, 2, \dots, Q$. This can be done by using (6) or the conclusion of Proposition 3. Now let $K = [-M, M]$. We see that for any $y = (x - \alpha n)/n^{1/m} > M$ or equivalently $x \notin \alpha_q n + K n^{1/m}$

$$\left| n^{-1/m} e^{-ix\xi_q} \hat{\phi}(\xi_q)^n H_m^{\beta_q}\left(\frac{x-\alpha_q n}{n^{1/m}}\right)\right| < \frac{Cn^{-1/m}}{2Q} \tag{65}$$

for all $q = 1, 2, \dots, Q$. Further, by combining (64) and (65) there is some natural number N such that

$$|\phi^{(n)}(x)| \leq C n^{-1/m}$$

for all $x \notin \cup_q (\alpha_q n + K n^{1/m})$ and $n > N$. Thus by Theorem 1, the supremum, $||\phi^{(n)}||_\infty$ must be attained on the set $\cup_q(\alpha_q n + K n^{1/m})$ for all $n > N$. Lastly, observe that by enlarging the compact set K, the above dependence on N can be removed. This completes the proof.

□

8 Examples

We begin this final section by considering a complex valued function on $\mathbb{Z}$ whose convolution powers consist of two waves drifting apart. This example is not able to be treated by the results of Schoenberg, Greville or Thomée.

Example 1. *Consider the function ϕ defined by*

$$\phi(0) = \frac{3}{8} \qquad \phi(\pm 2) = -\frac{1}{4} \qquad \phi(\pm 3) = \frac{i}{3} \qquad \phi(\pm 4) = \frac{1}{16}$$

and $\phi = 0$ otherwise. The convolution powers, $\phi^{(n)}$, exhibit two distinct wave packets drifting apart, each with a rate of $2n$ from $x = 0$. Figure 5 illustrates this behavior.



Figure 5: $|\phi^{(n)}|$ for $n = 50, 100$

The Fourier transform of ϕ is given by

$$\hat{\phi}(\xi) = \frac{3}{8} - \frac{1}{2}\cos(2\xi) + \frac{2i}{3}\cos(3\xi) + \frac{1}{8}\cos(4\xi).$$

Here, $\sup|\hat{\phi}| = 1$ *and is attained only at $\pi/2, -\pi/2 \in (-\pi, \pi]$. It follows that*

$$\log\left(\frac{\hat{\phi}(\xi \pm \pi/2)}{\hat{\phi}(\pm\pi/2)}\right) = \pm 2i\xi \mp \frac{5i}{3}\xi^3 - \frac{7}{3}\xi^4(1 + o(1)) \text{ as } \xi \to 0$$

and so by Theorem 3 (or Theorem 6)

$$n^{1/3}\phi^{(n)}(x) = e^{-ix\pi/2} H_3^{\frac{5i}{3}}\left(\frac{x - 2n}{n^{1/3}}\right) + e^{ix\pi/2} H_3^{\frac{-5i}{3}}\left(\frac{x + 2n}{n^{1/3}}\right) + o(1)$$

where the limit is uniform in x. This also can be written as

$$\begin{aligned}
\phi^{(n)}(x) &= n^{-1/3}(i)^x\left[H_3^{\frac{i}{3}}\left(-\frac{x+2n}{(5n)^{1/3}}\right) + (-1)^{x+1} H_3^{\frac{i}{3}}\left(\frac{x-2n}{(5n)^{1/3}}\right)\right] + o(n^{-1/3}) \\
&= n^{-1/3}(i)^x\left[Ai\left(-\frac{x+2n}{(5n)^{1/3}}\right) + (-1)^{x+1} Ai\left(\frac{x-2n}{(5n)^{1/3}}\right)\right] + o(n^{-1/3}) \\
&= f(n,x) + o(n^{-1/3}) \qquad (66)
\end{aligned}$$

where Ai denotes the standard Airy function. To appreciate the local limit theorems, 2 and 3, we consider $\phi^{(n)}(x)$ *for* $n = 10000$ *near the right wave packet* ($19700 \leq x \leq 20150$)*. Figure 6 shows the graph of* $\mathrm{Re}(\phi^{(n}(x))$ *and Figure 7 shows the approximation,* $f(n,x)$ *defined by* (66).



Figure 6: $\mathrm{Re}(\phi^{(n)})$ for $n = 10000$



Figure 7: $\mathrm{Re}(f(n,x))$ for $n = 10000$

What appears to be noise in Figure 7 is the oscillatory tail of the first term of the two comprising $f(n,x)$ *in* (66), $n^{-1/3}(i)^x Ai\left(-\frac{x+2n}{(5n)^{1/3}}\right)$*. Removing this* *term, we consider*

$$\begin{aligned} g(n,x) &= f(n,x) - n^{-1/3}(i)^x Ai\left(-\frac{x+2n}{(5n)^{1/3}}\right) \\ &= -n^{-1/3}(-i)^x Ai\left(\frac{x-2n}{(5n)^{1/3}}\right). \end{aligned}$$

As Figure 8 shows, $g(n,x)$ is a much better approximation of $\phi^{(n)}(x)$ at $n = 10000$ for $19700 \leq x \leq 20150$. As one can check, $g(n,x)$ is precisely the function yielded by Theorem 2.



Figure 8: $\mathrm{Re}(g(n,x))$ for $n = 10000$

Example 2. *We return to the example given in the introduction and justify the claims made therein. Let ϕ be given by (1). A quick computation shows that*

$$\hat{\phi}(\xi) = 1 - \frac{i}{2}\sin^2(\xi/2) - \sin^4(\xi/2)$$

where the supremum of $|\hat{\phi}|$ on the interval $(-\pi,\pi]$ is only attained at $\xi_0 = 0$. In the notation of Proposition 1 we may write

$$\eta(\xi) = \log\left(\frac{\hat{\phi}(\xi)}{\hat{\phi}(0)}\right) = -i\xi^2(\frac{1}{8} - \frac{1}{96}\xi^2) - \frac{7}{128}\xi^4 + \sum_{l=5}^{\infty} a_l \xi^l$$

on a neighborhood of 0 and so by the proposition, $m = 2$, $\alpha = 0$ and $\beta = i/8$. By Theorem 1, there are constants $C, C' > 0$ such that

$$Cn^{1/2} \leq ||\phi^{(n)}||_\infty \leq C'n^{1/2}. \tag{67}$$

By Theorem 2 and using (5) *we may also conclude that*

$$\begin{aligned}\phi^{(n)}(\lfloor xn^{1/2}\rfloor) &= n^{-1/2}H_2^{i/8}(x)+o(n^{-1/2})\\ &= \frac{n^{-1/2}}{\sqrt{4\pi i/8}}e^{-8|x|^2/4i}+o(n^{-1/2})\end{aligned}$$

where the limit is uniform for x in any compact set.

Example 3. *In the article [2], Example 2.4 and Proposition 2.5 therein described the asymptotic behavior of the convolution powers of an arbitrary real valued function ϕ supported on three (consecutive) points. In the notation of the proposition we define ϕ by*

$$\phi(0)=a_0,\ \phi(\pm 1)=a_\pm \text{ and } \phi=0 \text{ otherwise} \tag{68}$$

where $a_0, a_+, a_- \in \mathbb{R}$. As in [2] we will also assume that $a_0 > 0$ and that $a_+ \neq 0$ or $a_- \neq 0$; this assumption guarantees that ϕ has admissible support. Proposition 2.5 of [2] describes the asymptotic behavior of $\phi^{(n)}$ for all values of $a_0, a_\pm$ except the special case where $a_+a_- < 0$ and $4|a_+a_-| = a_0|a_+ + a_-|$. Theorem 3 of the present article allows us to treat this final case with ease. We do this now.

Proposition 5. *Let ϕ be as above and assume additionally that $a_+a_- < 0$ and $4|a_+a_-| = a_0|a_+ + a_-|$. If $a_+ + a_- > 0$ then*

$$\phi^{(n)}(x) = n^{-1/3}A^n H_3^{\beta}\left(\frac{x-\alpha n}{n^{1/3}}\right) + o(A^n n^{-1/3}) \tag{69}$$

uniformly for $x \in \mathbb{Z}$ where $A = a_0 + a_+ + a_-$, $\alpha = (a_+ - a_-)/A$ and $\beta = i(\alpha - \alpha^3)/6$.

If $a_+ + a_- < 0$ then

$$\phi^{(n)}(x) = n^{-1/3}A^n e^{-ix\pi} H_3^{\beta}\left(\frac{x-\alpha n}{n^{1/3}}\right) + o(A^n n^{-1/3}) \tag{70}$$

uniformly for $x \in \mathbb{Z}$ where $A = a_0 - a_+ - a_-$, $\alpha = (a_- - a_+)/A$ and $\beta = i(\alpha - \alpha^3)/6$.

In either case, there is a compact set K for which the $||\phi^{(n)}||_\infty$ is attained on the set $(\alpha n + K n^{1/3})$.

Proof. We may write

$$\hat{\phi}(\xi) = a_0 + a_+e^{i\xi} + a_-e^{-i\xi} = a_0 + (a_+ + a_-)\cos(\xi) + i(a_+ - a_-)\sin(\xi).$$

Under the assumption that $4|a_+a_-| = a_0|a_+ + a_-|$ and $a_+ + a_- > 0$, it was shown in [2] that $|\hat{\phi}|$ is maximized only at $0 = \xi_0 \in (-\pi, \pi]$ and in which case this maximum takes the value $A = a_0 + a_+ + a_-$.

Set $\psi(x) = \phi(x)/A$. It follows immediately that $A^n\psi^{(n)}(x) = \phi^{(n)}(x)$ and $\sup|\hat{\psi}| = 1$ which is taken only at $\xi_0 = 0$. In the notation of Proposition 1 we have

$$\eta(\xi) = \log\left(\frac{\hat{\psi}(\xi)}{\hat{\psi}(0)}\right) = i\left(\frac{(a_+ - a_-)}{a_0 + a_+ + a_-}\right)\xi$$
$$- \frac{i}{6}\left(\frac{(a_+ - a_-)(a_0^2 - a_0a_+ - a_0a_- - 8a_+a_-)}{(a_0 + a_+ + a_-)^3}\right)\xi^3 - \gamma\xi^4 + \sum_{l=5}^{\infty} a_l\xi^l$$

on a neighborhood of 0 where $\gamma > 0$. Setting $\alpha = (a_+ - a_-)/A$ and using the fact that $4|a_+a_-| = a_0|a_+ + a_-|$ we write

$$\eta(\xi) = i\alpha\xi - \frac{i}{6}(\alpha - \alpha^3)\xi^3 - C\xi^4 + \sum_{l=5}^{\infty} a_l\xi^l \tag{71}$$

on a neighborhood of 0. By a quick inspection of (71) it is clear that ψ meets the hypotheses of Theorem 3 where $m = 3$ and $\beta = i(\alpha - \alpha^3)/6$. Therefore

$$\psi^{(n)}(x) = n^{-1/3}H_3^{\beta}\left(\frac{x - \alpha n}{n^{1/3}}\right) + o(n^{-1/3}) \tag{72}$$

uniformly for $x \in \mathbb{Z}$. The limit (69) follows immediately by multiplying (72) by A^n. An appeal to (9) of Theorem 3 shows that $||\psi^{(n)}||_\infty$ and hence $||\phi^{(n)}||_\infty$ is indeed attained on the set $(\alpha n + Kn^{1/3})$ for some compact set K.

In the case that $a_+ + a_- < 0$ it was shown in [2] that $|\hat{\phi}|$ attains its only maximum at $\xi_0 = \pi \in (-\pi, \pi]$. Upon setting $A = a_0 - a_+ - a_-$, $\psi(x) = \phi(x)/A$ and considering the Taylor expansion of $\log(\hat{\psi}(\xi + \xi_0)/\hat{\psi}(\xi_0))$, the result follows by an argument similar to that given for the previous case. □

References

[1] P. Brenner, V. Thomée, and L. Wahlbin, Besov Spaces and Applications to Difference Methods for Initial Value Problems. Springer-Verlag, Berlin, 1975.

[2] P. Diaconis and L. Saloff-Coste: Convolution powers of complex functions on Z. arXiv:1205.6490.

[3] I. M. Gelfand and G. E. Shilov: Generalized Functions, II, Academic Press, New York and London, 1968.

[4] T.N.E. Greville: On Stability of Linear Smoothing Formulas. *SIAM J. Numer. Anal.,* 3(1):157-170, 1966.

[5] R. Hersh: A class of "Central Limit Theorems" For Convolution Products of Generalized Functions. *Trans. Amer. Math. Soc.,* 140:71-85, 1969.

[6] G. Lawler and V. Limic: Random Walk: A Modern Introduction. Cambridge University Press, New York, 2010.

[7] I.J. Schoenberg: On smoothing operations and their generalized functions. *Bull. Amer. Math. Soc.,* 59:299-230, 1953.

[8] F. Spitzer: Principle of Random Walk, 2nd ed. Springer-Verlag, New York, 1976.

[9] Vidar Thomée. Stability theory for partial difference operators. *SIAM Rev.*, 11:152-195, 1969.

[10] Vidar Thomée. Stability of difference schemes in the maximum-norm. *J. Differential Equations,* 1:273-292, 1965.